

Microbiological processes in banded iron formation deposition

NICOLE R. POSTH*¹, KURT O. KONHAUSER† and ANDREAS KAPPLER*

*Center for Applied Geosciences, University of Tuebingen, 72076, Tuebingen, Germany

(E-mail: andreas.kappler@uni-tuebingen.de)

†Department of Earth and Atmospheric Sciences, University of Alberta, Edmonton, Alberta, Canada T6G 2E3

ABSTRACT

Banded iron formations have been studied for decades, particularly regarding their potential as archives of the Precambrian environment. In spite of this effort, the mechanism of their deposition and, specifically, the role that microbes played in the precipitation of banded iron formation minerals, remains unresolved. Evidence of an anoxic Earth with only localized oxic areas until the Great Oxidation Event *ca* 2.45 to 2.32 Ga makes the investigation of O₂-independent mechanisms for banded iron formation deposition relevant. Recent studies have explored the long-standing proposition that Archean banded iron formations may have been formed, and diagenetically modified, by anaerobic microbial metabolisms. These efforts encompass a wide array of approaches including isotope, ecophysiological and phylogeny studies, molecular and mineral marker analysis, and sedimentological reconstructions. Herein, the current theories of microbial processes in banded iron formation mineral deposition with particular regard to the mechanisms of chemical sedimentation and post-depositional alteration are described. The main findings of recent years are summarized and compared here, and suggestions are made regarding cross-disciplinary information still required to constrain the role of the biosphere in banded iron formation deposition.

Keywords Aerobic and anaerobic Fe(II) oxidation, anoxygenic phototrophs, atmospheric evolution, banded iron formations, cell-mineral aggregates, mineral diagenesis.

INTRODUCTION

Precambrian banded iron formation (BIF) deposition continues to be a topic of research interest because their mineral and chemical compositions provide direct evidence of the environmental conditions that existed at the time of their deposition. In particular, a number of recent studies have focused on how BIFs can be used as proxies for the emergence of oxygen, from its initial production in the oceans by at least 2.7 Ga (Eigenbrode & Freeman, 2006; Godfrey & Falkowski, 2009; Kendall *et al.*,

2010), to its spread throughout the atmosphere between 2.45 Ga and 2.32 Ga, the so-called Great Oxidation Event (GOE) (Rye & Holland, 1998; Pavlov & Kasting, 2002; Bekker *et al.*, 2004; Buick, 2008; Farquhar *et al.*, 2011; Konhauser *et al.*, 2011a,b). In this regard, BIFs offer an archive of both early microbial life and the transition to an oxygen-dominated Earth system.

At the heart of understanding the formation of these deposits lies the potential interaction of abiotic and biological processes in these ancient systems; both abiotic and biotic models of formation exist and continue to be explored. The

¹Nordic Centre for Earth Evolution (NordCEE), Institute for Biology, University of Southern Denmark, 5230, Odense, Denmark

relevant question to the efficacy of the microbial models in terms of BIF precipitation is whether the dominant processes of Fe(II) oxidation and mineral precipitation can be deduced from their: (i) microfossils and fabrics; (ii) stable isotope composition; (iii) biosignature identity; (iv) geochemical composition; and (v) sedimentological features. Moreover, if a biological mechanism was important in the initial process of Fe(II) oxidation in the ancient ocean water column, it is expected that biomass would have settled to the sea floor along with the Fe(III) minerals. This organic carbon would subsequently have served as an oxidizable substrate for sedimentary microbial communities during diagenesis and metamorphism. These post-depositional processes would have modified the initial BIF precursor sediment into the mineral assemblages observed today. Herein, the existing evidence for microbiological involvement in both BIF precipitation and burial is summarized.

BANDED IRON FORMATION SETTING AND DEPOSITIONAL THEORIES

Banded iron formations are iron-rich (*ca* 20 to 40% Fe) and siliceous (*ca* 40 to 50% SiO₂) sedimentary deposits that precipitated throughout much of the late Archean (2.7 to 2.5 Ga) and Palaeoproterozoic (2.5 to 1.8 Ga). The 'Superior' type BIFs, including those in the Hamersley Group, Western Australia, and the Transvaal Supergroup, South Africa, are hundreds of metres thick, over 10⁵ km² in areal extent and contain >10¹³ tons of iron (Beukes, 1984; Trendall, 2002). These formations are characteristically laminated (James, 1954; Fig. 1), with banding observed over a wide range of scales, from coarse macrobands (metres in thickness) to mesobands (centimetre-thick units) to millimetre and sub-millimetre layers (Trendall & Blockley, 1970). Among the latter are the various varve-like repetitive laminae, known as microbands. Although it is not resolved that these microbands are seasonal, much of the work on these structures is based on that premise.

The mineralogy of the least metamorphosed BIFs consists of chert, magnetite, hematite, carbonates (siderite and dolomite-ankerite), greenalite, stilpnomelane and riebeckite (Klein, 2005; Bekker *et al.*, 2010); the presence of both ferric and ferrous minerals gives BIFs an average oxidation state of Fe^{2.4+} (Klein & Beukes, 1992). It is generally agreed that none of the minerals in BIFs are primary, but that instead, the mine-

rals reflect both diagenetic and metamorphic overprinting. For instance, the primary iron minerals were most probably ferric hydroxide [Fe(OH)₃], greenalite [(Fe)₃Si₂O₅(OH)₄] and siderite (FeCO₃) (Han, 1966, 1978; Dimroth & Chauvel, 1973; Perry *et al.*, 1973; Klein, 2005). It has also been suggested that the Archean ocean had significantly elevated concentrations of dissolved silica, at least as high as at saturation with cristobalite (0.67 mM at 40°C in sea water), and possibly even amorphous silica (2.20 mM) (Maliva *et al.*, 2005; Konhauser *et al.*, 2007a). Under such silica-rich conditions, the precipitation of amorphous silica could have taken place directly on the sea floor (Krapež *et al.*, 2003).

Although BIFs are generally believed to have been deposited on the continental shelf at water depths >200 m due to the lack of obvious current and wave-generated structures (Beukes, 1973; Klein & Beukes, 1992; Morris & Horwitz, 1983; for review see Klein, 2005), it has been argued that some were deposited off the continental slope in the deep ocean (Krapež *et al.*, 2003). The dominant source of Fe(II) into the Archean ocean was hydrothermal (Jacobsen & Pimentel-Klose, 1988; Bau & Möller, 1993; Hamade *et al.*, 2003), but the proximity of the Fe(II) source to the site of deposition is still unclear. It is believed that the concentration of Fe(II) in these basins ranged from 0.05 to 0.5 mM (Holland, 1973; Morris, 1993). On the one hand, Fe could have been delivered from the deep ocean to the outer continental shelf by upwelling currents from a mid-ocean ridge system (Holland, 1973; Morris & Horwitz, 1983). Accordingly, BIFs would sediment from below the wave base (without the influence of wave and storm-induced currents) onto partially submerged platforms of the continental shelves. On the other hand, the direct supply of Fe(II) into the photic zone by hydrothermal plumes associated with shallow seamount-type systems (Isley, 1995; Isley & Abbot, 1999) would curtail the difficulties introduced by the high upwelling rates needed to bring sufficient iron from the deep sea onto the continental shelf (Konhauser, 2007b).

The mineralogy of BIFs dictates that some oxidation of Fe(II) was necessary for formation, although the dominant mechanism(s) is uncertain (Fig. 2). Prior to the rise of atmospheric oxygen and the development of a protective ozone layer, the Earth's surface was subjected to high levels of ultraviolet (UV) radiation. Bulk ocean waters that were anoxic at this time could have supported high concentrations of dissolved Fe(II). Under





Fig. 1. Overview of the laminated BIF from the Joffre iron formation, Pilbara Craton, north-west Australia. Image courtesy of Stefan Lalonde (A). Core consists of red microbands (<1 mm) of chert-hematite-riebeckite (bluish bands in upper core) alternating with lighter chert–dolomite–siderite–crocidolite mesobands (≥ 1 cm) and denser, dark magnetite mesobands. A slumped chert band due to post-depositional compaction is visible in core top (B). A scanned thin section of the Joffre iron formation in plane-polarized light (C) shows dense blue riebeckite microbands interbedded with brownish hematite/goethite microbands and chert microbands and mesobands. The chert mesobands contain wavy, fine-grained carbonate and hematite/goethite microbands (image scale: 2.6×4.6 cm). Images (B) and (C) courtesy of Rasmus Haugaard.

such conditions, dissolved ferrous iron species, such as Fe^{2+} or $\text{Fe}(\text{OH})^+$, would have absorbed radiation in the 200 to 400 nm range, leading to the formation of dissolved ferric iron [reaction 1], which, in turn, hydrolysed to form ferric hydroxide at circumneutral pH (Cairns-Smith, 1978; Braterman *et al.*, 1983; Fig. 2A).



However, these experiments focused on determining the specific rates of Fe(II) photochemical oxidation, and did not simulate the complex, disequilibrium water chemistry characteristic of an ocean where Fe(II)-rich hydrothermal waters reacted with ambient Si-saturated sea water that also contained high concentrations of HCO_3^- . Indeed, in fluids with high dissolved Fe(II), $\text{Si}(\text{OH})_4$ and HCO_3^- , the oxidation effects of

either UVA or UVC were found to be negligible compared with the precipitation of ferrous-iron silicates (Konhauser *et al.*, 2007b).

As an alternative to the abiological model, the presence of ferric iron minerals in BIFs has also been ascribed to the metabolic activity of planktonic bacteria in the ocean's photic zone. Chemical oxidation of Fe(II) by photosynthetically produced O_2 is one possibility, allowing for the indirect biogenic precipitation of ferric hydroxide (Fig. 2B and C). Under an anoxic atmosphere, this O_2 could have been confined to localized 'oxygen oases' associated with cyanobacterial blooms in coastal settings (Cloud, 1965, 1973). Cloud (1965, 1973) further proposed that such primitive O_2 -producing photosynthetic bacteria, which lacked suitably advanced oxygen-mediating enzymes, required ferrous iron to detoxify oxygen. If so, these micro-organisms would have




Fig. 2. Models of BIF deposition: For an anoxic water column, the proposed abiotic mechanism of Fe(II) photooxidation by UV light (A). The traditional model of BIF deposition involves production of oxygen by cyanobacteria, which then chemically reacts with hydrothermal dissolved Fe(II). The restriction of cyanobacterial mats to the near-shore would physically separate Fe(III) oxyhydroxides from organic carbon precipitates (B), which would not be the case in a system where cyanobacteria also populated off-shore regions (C). A biotic mechanism of deposition proposed for an anoxic setting is the direct microbial oxidation of Fe(II) via anoxygenic Fe(II)-oxidizing phototrophy (D).

flourished specifically when Fe(II) and nutrients were made episodically available (Cloud, 1969; Fralick & Pufahl, 2006). Once oxygen was present, microaerophilic Fe(II) oxidizers could have contributed to the direct precipitation of ferric hydroxide (Holm, 1989).

Garrels *et al.* (1973) and Hartman (1984) later suggested that light, not O₂, may have coupled the carbon and iron cycles, via photosynthesis that used Fe(II) rather than H₂O as an electron donor, producing Fe(III) rather than O₂ (Fig. 2D). With the evolution of this photoferrotrophy, biological Fe(II) oxidation would have superseded photochemical oxidation because the bacteria could grow deeper in the water column where UV radiation would be effectively absorbed (Kohnhauser *et al.*, 2002). Modelling of experimentally determined photosynthetic Fe(II) oxidation rates even suggests that such micro-organisms could have accounted for all of the Fe(III) initially deposited in primary BIF sediment (Kappler *et al.*, 2005).

Although the average oxidation state of the iron in BIFs (Fe^{2.4+}) can be attributed to the simultaneous deposition of Fe(II) and Fe(III), the partial microbial or abiotic reduction of Fe(III) to

Fe(II) can also explain this mixed oxidation state. The presence of magnetite as: (i) disseminated grains within but obscuring sedimentary laminae; (ii) laminated beds that clearly truncate sedimentary layering; and (iii) cleavage fills, suggests a secondary origin to magnetite in BIFs (Ewers & Morris, 1981; Krapež *et al.*, 2003). Microbial Fe(III) reduction is carried out by anaerobic heterotrophic bacteria that link this process to organic matter oxidation (Lovley & Phillips, 1988). Fe(III) oxyhydroxides can also be abiotically reduced after sedimentation with a reductant, such as the organic carbon associated with planktonic bacteria necromass, and can produce siderite (FeCO₃) and magnetite (Fe₃O₄) (Schaefer *et al.*, 2009; Posth *et al.*, 2010). Recent experiments point to the production of siderite spherulites, also found in many BIFs, by the experimental diagenesis of ferrihydrite and glucose at relevant pressure and temperature. Three lines of evidence point to the antiquity of such an anaerobic respiratory pathway. Firstly, extant hyperthermophilic *Bacteria* and *Archaea* branch deeply in the universal phylogenetic tree and can reduce Fe(III) (Vargas *et al.*, 1998). Secondly, highly negative $\delta^{56}\text{Fe}$ values in magnetite-rich

BIF samples as old as 2.9 Ga have been observed (Johnson *et al.*, 2003; Yamaguchi *et al.*, 2005) and are comparable to the negative fractionations measured in experimental culture with dissimilatory Fe(III)-reducing bacteria (Johnson *et al.*, 2005). Thirdly, Li *et al.* (2011) recently conducted high-resolution mineral analyses of the 2.48 Ga Dales Gorge BIF in Western Australia and reported the detection of an Fe(III)-acetate salt, as well as nanocrystals of apatite in association with magnetite; the latter had a crystal chemistry identical to modern biogenic magnetite. This combination of features points to the original presence of biomass in the BIF sediments, and also indicates that the organic carbon served as an electron donor during bacterial Fe(III) reduction.

Crucial to understanding of the biological role in BIF formation is an assessment of the relation between biomass production and the BIF depositional setting. To this end, results of physiological experiments give a sense of the viable range of initial phototrophic productivity in the ocean's photic zone and of precipitated cell-Fe(OH)₃ aggregate composition. Investigation of the redox properties and the cycling of elements have also proven valuable in the reconstruction of the microbial role at the time of BIF deposition. The utilization of H₂O by cyanobacteria and the direct oxidation of Fe(II) by photoferrotrophs are both coupled to CO₂ reduction and biomass production. Ferric hydroxide particles and organic biomass could be deposited in either case. In the photoferrotroph model, direct Fe(II) oxidation by the cells would lead to a joint sedimentation of precipitated Fe(III) and microbially produced biomass. In the cyanobacteria model, the production of O₂ by the cells, and the oxidation of the Fe(II) by this O₂ are independent. Consequently, the potential association of the Fe(III) minerals with the cyanobacterial biomass depends on the local distribution of the cells relative to the location of Fe(III) mineral deposition (open ocean or shore, Fig. 2C).

EVIDENCE FOR ANCIENT CYANOBACTERIA

Microfossils and biological fabrics

The timing of the origins of cyanobacteria remains highly controversial. For many years, the oldest morphological evidence for cyanobacteria was believed to have come from weathered clasts in the 3.45 Ga Apex cherts of the

Warrawoona Group, Western Australia. Structures reminiscent of unbranched, partitioned trichomes, with dimensions and morphologies indicative of extant cyanobacteria, led to the suggestion that the Archean micro-organisms were capable of gliding and, possibly, phototactic motility (Schopf, 1993). Moreover, confocal laser scanning microscopy of the filaments and Raman spectroscopy were used to infer the presence of biogenic kerogen in higher concentrations than the surrounding matrix (Schopf *et al.*, 2002; Schopf & Kudryavtsev, 2009). Additional support for the early evolution of cyanobacteria came from the discovery of large spheroidal, sheath-like structures (up to 20 µm in diameter) in cherts from the underlying Towers Formation (Schopf & Packer, 1987), as well as from the presence of conical and pseudocolumnar stromatolites in the 3.4 Ga Strelley Pool Chert, Western Australia (Van Kranendonk, 2006).

A re-examination of the Apex chert by Brasier *et al.* (2002, 2005), however, called into question the biogenicity of the filamentous structures and the sedimentary origins of the earliest 'fossiliferous' deposits, citing Fischer-Tropsch-type reactions associated with sea floor hydrothermal systems. Buick (1988) has maintained that the actual chert units from which these microfossils derive are secondary hydrothermal deposits of much younger age that cross-cut the primary bedding. The Warrawoona samples also led to experiments which demonstrate that kerogen in a microfossiliferous sample does not in itself indicate biogenicity. Simple organic hydrocarbons, whose sources are abiogenic (formaldehyde), readily condense onto silica-carbonate inorganic filaments and subsequently polymerize under gentle heating to yield kerogenous products (Garcia-Ruiz *et al.*, 2003). Indeed, the same kind of Raman spectral signature is obtained from kerogen as stems from many other poorly ordered and abiogenic carbonaceous materials (Pasteris & Wopenka, 2003). Furthermore, a recent study focusing on similar structures from the Apex chert using Raman spectroscopy showed that the microstructures were a mixture of hematite and quartz and that the matrix did in fact consist of carbonaceous material, but the interpretation of the original structures as biogenic was a result of image compilation (Marshall *et al.*, 2011). Thus, the discussion surrounding the Warrawoona samples demonstrates that the presence of carbonaceous material is not sufficient proof of biogenicity. Importantly, it highlights the stringent criteria

needed to determine *bona fide* biogenicity in ancient rocks (for further discussion on these criteria, see Schopf *et al.*, 2007). A firmer benchmark for the presence of cyanobacteria comes from the 2.7 Ga stromatolitic assemblages of the Tumbiana Formation, Western Australia. Based on their habitat in a sulphate-deficient evaporative lake, it was suggested that the original microbial mat contained cells that were metabolized by oxygenic photosynthesis (Buick, 1992). This view is supported by the earliest recognized fossil assemblage of filamentous and coccoid cells colonies, from the 2.6 Ga Campbell Group, South Africa, which appears to include oscillatoriacean cyanobacterial genera, such as *Phormidium* or *Lyngbya*. These micro-organisms contributed to the formation of stromatolitic reefs in shallow subtidal to intertidal settings (Altermann & Schopf, 1995). Other possible examples of ancient cyanobacteria occur in laminates and oolites of the 2.5 Ga Gamohaan Formation, South Africa, where Wright & Altermann (2000) reported dolomicrites formed on the outer margins of the sheaths of filamentous cyanobacterial remains. More putative evidence for cyanobacteria stems from the 2.9 Ga Pongola Supergroup in South Africa, where Noffke *et al.* (2008) observed microbially influenced sedimentary structures that are consistent with similar features constructed today by benthic cyanobacteria.

The link between oxygenic photosynthesis and stromatolite formation is supported by modern studies which show that the trapping and binding of carbonate grains by cyanobacterial filaments and their extracellular polymers (such as extracellular polymeric substances) are integral to the structure of intertidal stromatolites (Burns *et al.*, 2004, 2009; Bosak *et al.*, 2009). However, a series of studies show how similar structures can be formed via abiological processes (Grotzinger & Rothman, 1996; Grotzinger & Knoll, 1999; McLoughlin *et al.*, 2008). In addition, sequence analysis of small sub-unit rRNA genes amplified with PCR from genomic DNA of modern stromatolite communities of Hamelin Pool, Shark Bay, Australia, showed that anoxygenic phototrophs represent a considerable fraction of the biomass (Papineau *et al.*, 2005). Along the same lines, Bosak *et al.* (2007) later demonstrated that the anoxygenic phototroph, *Rhodospseudomonas palustris*, stimulates the precipitation of calcite in saturated solutions and builds stromatolite-like structures. It has been suggested that before cyanobacteria evolved, ferrous iron was the

main reductant in ancient oceans; only with depletion in ferrous iron supply to the oceans did H₂O₂ and then water replace ferrous iron as the electron donor in the cyanobacterial line of evolution (Olson & Blankenship, 2004). It seems possible that prior to the rise of cyanobacteria, predecessor anaerobes, such as anoxygenic phototrophs, were dominant stromatolite-building organisms, which could account for the stromatolitic record that pre-dates strong evidence for oxygenic phototrophs. Further studies into the mechanisms of stromatolite formation promise to help parse the dominant processes to use these structures more accurately in ecosystem reconstruction.

Stable isotope composition

One persuasive argument in favour of early CO₂ fixation via photosynthesis, and the potential presence of cyanobacteria, comes from the highly negative carbon isotopic values of kerogen residues in Archean strata. The negative values are seemingly indicative of life because the transformation of inorganic carbon (for example, CO₂ or HCO₃⁻) via autotrophic pathways into organic carbon involves the preferential incorporation of the lighter isotope, ¹²C, into the organic material, leaving behind a reservoir enriched in the heavier isotope, ¹³C. Consequently, organic compounds produced by autotrophic pathways display a marked preference for the light isotope (for example, δ¹³C in cyanobacteria range from -4 to -35‰), whereas the heavy carbon is retained in the surface reservoir of oxidized carbon, mostly as dissolved bicarbonate, and later incorporated into precipitated carbonate minerals, for example, calcite and dolomite, or as atmospheric CO₂ (Schidlowski, 2000).

Analysis of kerogens in microfossiliferous units in Archean formations from Western Australia and South Africa contain negative carbon isotopic values that range from -20 to -35‰ (Altermann & Kazmierczak, 2003; Tice & Lowe, 2004), whereas the organic carbon in early Proterozoic cherts similarly averages *ca* -31.0 ± 4.7 (Strauss *et al.*, 1992). More recently, a study of the C-isotope signatures in the Transvaal Supergroup showed the organic δ¹³C to range from -40 to -25‰ (Fischer *et al.*, 2009). Collectively, these results are an indication of early photosynthesis. Unfortunately, a number of carbon fixation pathways (Calvin cycle and the reductive acetyl-CoA, reductive citric acid and hydroxypropionate pathways) utilized by phototrophic

bacteria seem to have overlapping degrees of carbon isotope fractionation, which hinders conclusive interpretation.

Strong evidence for oxygen in the Archean oceans comes from extremely isotopically depleted kerogens from 2.72 to 2.59 Ga carbonates and shales in the Hamersley Province in Western Australia. Organic carbon ^{13}C values range from as low as -57 to -28‰ (Eigenbrode & Freeman, 2006). The ^{13}C enrichment has been ascribed to the assimilation of methane by methanotrophic bacteria that utilize electron acceptors such as O_2 , SO_4^{2-} or NO_3^- ; both sulphate and nitrate require oxygen for their formation (Hayes, 1983). Potentially, although not yet conclusively demonstrated, methane oxidation could be linked to the reduction of Fe(III) (Konhauser *et al.*, 2005). Interestingly, there is a ^{13}C enrichment of 10‰ in kerogen in post-2.7 Ga shallow-water facies relative to the deep-water facies, suggesting that the shallow setting became oxygenated, marginalizing the methanogens and methanotrophs to deeper waters, whereas the shallow waters became dominated by cyanobacteria (Eigenbrode & Freeman, 2006).

Biomarker identity

An additional tool for the identification of a specific group of organisms in ancient rocks is the analysis of organic biomarkers. These organic compounds, derived from more complex precursors, retain some resemblance to the original biological molecules, even after a long history of decomposition and alteration that accompanies burial and diagenesis. Bitumens from the 2.6 Ga Marra Mamba Iron Formation and the 2.5 Ga Mount McRae Shale of the Hamersley Group, Western Australia, for example yield abundant 2α -methylhopanes, derivatives of prominent lipids in cyanobacteria (methyl-bacteriohopanepolyols) that serve to improve cell membrane rigidity (Brocks *et al.*, 1999; Summons *et al.*, 1999; Eigenbrode *et al.*, 2008).

These compounds were first strictly found in cyanobacteria and, therefore, were interpreted as diagnostic of oxygenic photosynthetic organisms in ancient rocks. The belief that they only served a structural function, however, made it difficult to connect the presence of these compounds to a specific metabolism. Genomic databases revealed that several facultative and obligate anaerobes possess the appropriate genes for hopanoid biosynthesis. In fact, *Geobacter sulfurreducens* actually produces a wide variety

of complex hopanoids structurally related to 2α -methylhopane under strictly anoxic conditions in pure culture (Fischer *et al.*, 2005; Härtner *et al.*, 2005). It has been demonstrated that an anoxygenic Fe(II)-oxidizing phototroph, *R. palustris* strain TIE-1, generates substantial quantities of 2-methylhopanoids in the absence of oxygen (Rashby *et al.*, 2007). Utilizing this same strain, a recent study now shows that C-2 methylated hopanoids are preferentially synthesized by the *R. palustris* strain TIE-1 under conditions of elevated Fe(II) concentrations (Eickhoff *et al.*, 2013). Accordingly, their presence in the rock record may also signal anoxic ferrous conditions. Although the full physiological roles of 2-methylhopanoids are still not defined, it seems that their origins are more diverse than first believed.

Hopanes carrying 3-methyl substituents have also been recovered from 2.72 to 2.56 Ga carbonates and shales from the Hamersley Province (Eigenbrode *et al.*, 2008). The only known extant bacteria to produce these particular hopanes are aerobic methanotrophs. Their presence, and strong correlation with the carbon isotope compositions in kerogen (as discussed in the previous section), may thus confirm the presence of oxygen in the photic zone by the late Archean.

A third suite of biomarkers, specific steranes of 28 to 30 carbon isomers, are unique alteration products of the sterols used in extant eukaryotic cell membranes. The only prokaryotes known to synthesize sterols have biosynthetic pathways leading to different structural isomers. Because O_2 is required for the biosynthesis of sterols, their extraction from Archean rocks suggests that at least some dissolved oxygen (*ca* $0.002 \text{ ml O}_2 \text{ l}^{-1}$) was present at that time. These steranes were reported in bitumens of the 2.7 Ga shales of the Jeerinah Formation, Hamersley Group (Brocks *et al.*, 1999). This finding was later questioned as a subsequent study suggested that the samples were contaminated by younger fluids (Rasmussen *et al.*, 2008). In recent experiments using the yeast *Saccharomyces cerevisiae* as a test organism, steroid biosynthesis was observed at dissolved O_2 concentrations ranging from $6.5 \mu\text{M}$ to 7 nM (Waldbauer *et al.*, 2011), suggesting that very low O_2 concentrations were sufficient for steroid biosynthesis. There is still no experimental evidence to support the alternative theory that steranes may have been produced via an anaerobic pathway utilizing hydrogen peroxide or organic peroxides (Raymond & Blankenship, 2004; Fischer & Pearson, 2007).

Geochemical composition

At present, there is a temporal gap in the rock record with regard to the geochemical evidence for the evolution of cyanobacteria and the effects of their metabolism on atmospheric oxygenation. In addition to the features described above that support cyanobacterial evolution *ca* 2.7 Ga, there are geochemical signatures in various lithologies that point towards the presence of shallow oxygenated sea water by the Neoproterozoic. Firstly, nitrogen isotopic composition of kerogens in minimally altered shales from the Campbellrand-Malmani platform in South Africa showed a significant rise in the $\delta^{15}\text{N}$ values by 2.67 Ga (Godfrey & Falkowski, 2009). This increase has been proposed as evidence for the onset of nitrification–denitrification reactions in the surface oceans; importantly, these microbial processes require the presence of oxygen. Secondly, a recent study of the trace element composition of those same shales showed a high abundance of Re but a low abundance of Mo, which together with the speciation of sedimentary iron, confirms the presence of oxygen in the water column, arguably to depths of several hundred metres (Kendall *et al.*, 2010). In contrast to the above, a number of features suggest that the atmosphere did not contain significant amounts of oxygen until *ca* 200 Myr later at *ca* 2.5 Ga, the so-called Great Oxidation Event. These include:

1 Detrital uraninite, pyrite and siderite that are easily oxidized by O_2 have been recovered in fluvial siliciclastic sediments of the 3.2 to 2.7 Gyr Pilbara craton (Rasmussen & Buick, 1999). They are not abundant in fluvial systems younger than 2.3 Ga.

2 High-resolution chemostratigraphy reveals an episode of enrichment of the redox-sensitive transition metals molybdenum and rhenium in the 2.5 Ga Mount McRae Shale in Western Australia (Anbar *et al.*, 2007). These findings suggest that the metals were supplied to the oceans by oxidative weathering of crustal sulphide minerals. Similarly, a recent compilation of Cr concentrations in BIFs shows an enrichment beginning at 2.45 Ga (Frei *et al.*, 2009; Konhauser *et al.*, 2011b). Given the insolubility of Cr minerals, its mobilization and incorporation into BIFs indicates enhanced chemical weathering at that time, likely associated with the evolution of aerobic continental pyrite oxidation.

3 Iron speciation and sulphur isotope data from the Mount McRae Shale provide evidence

for euxinic (anoxic and sulphidic) layers in the water column by 2.5 Ga (Reinhard *et al.*, 2009). These conditions were probably stimulated by an increase in oceanic sulphate concentrations arising from aerobic continental pyrite oxidation. Correspondingly, previous low levels of marine sulphate suggest minimal oxidative weathering of sulphide minerals (Canfield *et al.*, 2000).

4 Mass-independent fractionation (MIF) of sulphur isotopes ($\Delta^{33}\text{S}$ and $\Delta^{36}\text{S}$) in sulphide and sulphate-containing rocks deposited prior to 2.45 Ga, but not after 2.32 Ga. This observation indicates that the sulphur cycle changed from one governed by gas-phase photochemical reactions in an O_2 -deficient atmosphere to one dominated by oxidative weathering, where the different sulphur species lost their MIF signal due to microbial SO_4^{2-} reduction (Farquhar *et al.*, 2000; Mojzsis *et al.*, 2003; Bekker *et al.*, 2004). To preserve the MIF in Archean and early Palaeoproterozoic sediments, the atmospheric oxygen concentration must have been $<10^{-5}$ present atmospheric levels (PAL). By contrast, in atmospheres with O_2 concentrations $>10^{-5}$ PAL, reduced sulphur-bearing species would typically have been oxidized to sulphate before becoming incorporated into the sediment, so any MIF signature would have been lost (Pavlov & Kasting, 2002). A more recent and comparative study utilizing investigations of the Mount McRae Shale and correlating it to the Gamohaaran and Kuruman iron formations of South Africa dates the progressive rise of oxygen at 2.5 Ga and implies that this phenomenon was certainly on a widespread, if not global, scale (Kaufman *et al.*, 2007).

EVIDENCE FOR PHOTOFERROTROPHS

Given that pre-2.7 Ga BIFs probably were precipitated from anoxic sea water, and the various uncertainties regarding the presence of cyanobacteria in the rock record, it is relevant to consider BIF precipitation mechanisms based on the direct biological oxidation of Fe(II) via anoxygenic photosynthesis (Cloud, 1965, 1973; Garrels *et al.*, 1973; Hartman, 1984). Although there is still no actual physical or chemical evidence for the existence of Fe(II)-oxidizing phototrophs in the Archean, six independent lines do suggest their presence on the early Earth.


Phylogenetic analysis

Of the seven known strains of anoxygenic Fe(II)-oxidizing phototrophs known to date, six of them have been classified as *Proteobacteria*; the seventh is a green sulphur bacterium (Table 1). The *Proteobacteria* are a large and diverse phyla of bacteria consisting of five major classes; alpha [including Fe(II)-oxidizing purple non-sulphur bacteria], beta, gamma [including Fe(II)-oxidizing purple sulphur bacteria], delta and epsilon, all of which may have diversified from one ancestral phototroph (Woese, 1987). Purple sulphur and purple non-sulphur bacteria use bacteriochlorophyll (Bchl *a* or *b*) to absorb light in the near-infrared (800 to 1040 nm). Although controversy has arisen from phylogenetic studies of anoxygenic and oxygenic photosynthesis due to partial gene loss and fast lateral transfer of Bchl biosynthesis genes, molecular phylogenetic analyses of a number of enzymes involved in Bchl biosynthesis suggests that anoxygenic photosynthetic lineages are almost certain to be more deeply rooted than the oxygenic cyanobacterial lineages (Xiong, 2006).

Ecophysiological studies with extant species of anoxygenic Fe(II)-oxidizing phototrophs

Although the plausibility of an anoxygenic Fe(II)-oxidizing metabolism had long been speculated based on thermodynamic considerations (Garrels *et al.*, 1973; Hartman, 1984), such organisms were discovered more recently and subsequently isolated (Widdel *et al.*, 1993). A number of experimental studies have since confirmed that various purple and green phototrophic bacteria, both fresh water and marine (Table 1), can use Fe(II) as a reductant for CO₂ fixation [reaction 2] (Widdel *et al.*, 1993; Heising *et al.*, 1999; Straub

et al., 1999). This fact has made it possible to study these organisms as potential modern analogues of ancient organisms.



Anoxygenic Fe(II) oxidizing phototrophs are diverse and broadly distributed in the environment (Table 1). None of these organisms are unique in form as all are rod-shaped (Fig. 3); however, morphological variety is apparent, for example, the presence of vacuoles in *Thiodictyon* sp. strain F4 (Croal *et al.*, 2004). Furthermore, these strains are capable of utilizing a variety of substrates, such as acetate, H₂ and even FeS, and were shown to oxidize Fe(II) even in the presence of H₂ (Croal *et al.*, 2009). This diversity suggests that these organisms have the potential to be ubiquitous in both ancient and modern anoxic, Fe-rich environments.

Importantly, although the Fe(II) oxidation rate of these strains is dependent on light intensity, anoxygenic phototrophs are capable of oxidizing Fe(II) in low light regimes befitting the photic zone of ocean water (Kappler *et al.*, 2005; Hegler *et al.*, 2008). Considering the highly efficient light harvesting mechanism employed by anoxygenic phototrophs, it was estimated that, even in the presence of an overlying layer of cyanobacteria, sufficient light for the metabolism of anoxygenic phototrophs could penetrate up to 100 m ocean depth (Kappler *et al.*, 2005). Similar light harvesting has been seen in the Black Sea, with anoxygenic phototrophs detected at a depth of 80 to 100 m (Overmann *et al.*, 1992; Manske *et al.*, 2005). These strains also continue to carry out the oxidation of Fe(II) in the presence of a wide range of Fe(II) concentrations, with

Table 1. The anoxygenic Fe(II)-oxidizing phototrophs isolated to date include both marine and fresh water strains.

Phylum	Strain	Setting/location
α -Proteobacteria (purple non-sulphur)	<i>Rhodobacter ferrooxidans</i> sp. strain SW2	Fresh water ponds, Hannover Germany ¹
	<i>Rhodopseudomonas palustris</i> strain TIE-1	Fe-rich mat, Woods Hole, MA, USA ⁶
	<i>Rhodomicrobium vannielii</i> strain BS-1	Fresh water ditch, Tübingen, Germany ²
	<i>Rhodovulum iodosum</i>	Marine mud flat, North Sea, Germany ⁴
	<i>Rhodovulum robiginosum</i>	Marine mud flat, North Sea, Germany ⁴
γ -Proteobacteria (purple sulphur)	<i>Thiodictyon</i> sp. strain F4	Fresh water marsh, North Sea, Germany ⁵
Green sulphur	<i>Chlorobium ferrooxidans</i> sp. strain KoFox	Fresh water ditch, Constance, Germany ³

¹Ehrenreich & Widdel (1994), ²Heising & Schink (1998), ³Heising *et al.* (1999), ⁴Straub *et al.* (1999), ⁵Croal *et al.* (2004), ⁶Jiao *et al.* (2005).




Fig. 3. Anoxygenic Fe(II)-oxidizing phototrophic *Rhodospirillum rubrum* sp. strain SW2 cells form aggregates with the Fe(III) oxyhydroxides produced by their metabolism. Many cells are not encrusted by, but rather are associated with, the Fe(III) minerals (A), or are only partially coated (B).

reported values between 0.5 mM and 30 mM (Hegler *et al.*, 2008). This observation suggests that, even in ancient sea water with variations of Fe(II) concentrations due to a pulsed input of hydrothermal Fe(II), anoxygenic phototrophs could have continued to metabolize.

It has also been shown that Fe(II) oxidation is not hindered by the high concentrations of silica (2 mM) estimated for the Archean oceans (Siever, 1992; Konhauser *et al.*, 2007a). Indeed, the temperature dependence of anoxygenic Fe(II) oxidation in growth medium containing dissolved Fe(II) and dissolved silica suggests that a coupled abiotic–biotic mechanism may be involved in deposition of the alternating Fe-rich and Si-rich BIF layering (Posth *et al.*, 2008). Although demonstrated in a simplified laboratory system, these experiments show that, at temperatures suitable for Fe(II)-oxidizing phototrophs, the Fe(III) oxyhydroxides they produce will precipitate, whereas silica largely stays in solution. At cooler temperatures, silica will precipitate out abiotically whereas the bacteria cease measurable Fe(II) oxidation until temperatures once again increase.

Availability of substrates

Modern anoxygenic phototrophs are able to utilize multiple substrates. In Archean oceans, phototrophic bacteria would not have had access to large quantities of sulphide as an electron donor because any hydrothermally sourced dissolved sulphide would have reacted with Fe(II) near the vent and thus precipitated as solid-phase sulphide minerals. In the case of limited HS[−] supply, the ability of bacteria to use

Fe(II) as a reductant is predictable. In fact, even considering higher hydrogen concentrations (Kump & Barley, 2007), Fe(II) oxidation by modern analogue anoxygenic phototrophs still proceeds at significant rates under an atmosphere containing approximately three times the maximum predicted concentration of H₂ in the Archean atmosphere (Croal *et al.*, 2009). The input of dissolved Fe(II) from mid-ocean ridges was almost certainly greater during the Archean, a view supported by the presence of excess Fe in sandstones and shales of that time (Kump & Holland, 1992). Thus, it seems likely that these organisms applied enzymatic systems to use abundantly available electron donors, such as Fe²⁺ [reaction 2]. In addition, it was suggested that Fe(II) oxidation and Fe(III) precipitation may even have provided an external UV protecting shield for planktonic cells (Pierson, 1994; Phoenix *et al.*, 2001).

Biomarker identity

Recent analysis of the 1.64 Ga Barney Creek Formation in northern Australia, a marine, subtidal unit, yielded okenane, the fossil hydrocarbon biomarker of the precursor, okenone. This carotenoid pigment is exclusively found in purple sulphur bacteria and in recent sediments under euxinic conditions (Brocks, 2005). The presence of these biomarkers, therefore, strongly suggests that purple sulphur bacteria [that have representatives that are able to oxidize Fe(II) phototrophically] were present in high numbers, at least in the Palaeoproterozoic oceans. In addition, the discovery of 2-methylhopanoids in cultures of the anoxygenic Fe(II)-oxidizing

phototroph, *R. palustris* strain TIE-1 stimulated by environmental Fe(II) in the absence of oxygen (Rashby *et al.*, 2007) demonstrates the utility of this biomarker in reconstructing anoxygenic phototrophy in ancient sediment.

Stable isotope composition

Large variations of $\delta^{56}\text{Fe}$ (-2.5 to 1.0% relative to the bulk Earth) in late Archean and early Palaeoproterozoic BIFs from the Transvaal Supergroup, South Africa, incorporate the entire range of values measured on Earth and are interpreted to highlight the role of fluid-mineral isotope fractionation and, possibly, microbial processes (Johnson *et al.*, 2003). Although Fe isotope variations have the potential to define ancient sedimentary environments and could help distinguish between biotic and abiotic processes, as well as whether post-depositional processes may have altered the Fe redox state in BIFs, the complexity of the depositional environments and infiltration of external fluids during diagenesis and/or low temperature metamorphism may erase the primary isotope record (Hoefs, 1997; Johnson *et al.*, 2003). Yet, the Fe isotope composition of iron minerals in BIFs are markedly different from the homogenous values seen in igneous rocks and many modern marine sediments ($\delta^{56}\text{Fe}$ $0.00 \pm 0.05\%$) and the Fe isotope compositions of the major minerals found in BIFs (hematite, magnetite, Fe-carbonate and pyrite) reflect equilibrium fractionation of minerals, a variation in the fluid isotope composition from which the minerals precipitated, and microbial processes (Johnson *et al.*, 2003, 2004; Johnson & Beard, 2005; Steinhofel *et al.*, 2010).

A recent study focused on the rare earth element and Fe isotope archive in the iron and microfossil-rich stromatolites of the 1.89 Ga Gunflint and Biwabik iron formations (USA) with the aim of determining the dominant microbes involved in Fe(II) oxidation (Planavsky *et al.*, 2009). These authors point to the predominantly positive $\delta^{56}\text{Fe}$ values in this formation (ranging from -0.66 to $+0.82\%$) as evidence of Fe(II)-oxidizing bacteria, whereas Ce anomalies in these same formations suggest low oxygen conditions during deposition. As a comparison, laboratory experiments designed to define isotopic fractionation by anoxygenic Fe(II)-oxidizing phototrophs in pure culture, as well as enrichment cultures, show that ferric hydroxides are enriched in the heavy isotope by $1.5 \pm 0.2\%$ relative to Fe(II) (Croal *et al.*, 2004). This frac-

tionation was independent of the rate of the Fe(II) oxidation by these organisms. Heavier Fe(III) oxyhydroxides were also shown to be produced by microbially catalysed aerobic Fe(II) oxidation at low pH (Balci *et al.*, 2006), nitrate-reducing anaerobic Fe(II) oxidizers (Kappler *et al.*, 2010), aerobic chemical oxidation (Bullen *et al.*, 2001) and UV Fe(II) oxidation (Straton *et al.*, 2006), suggesting that all major processes produce similar Fe isotope fractionations. On the one hand, the isotopic fractionations reported from modern experiments correlate well with values measured in Archean to early Proterozoic BIFs from the Transvaal Supergroup, South Africa; they have $^{56}\text{Fe}/^{54}\text{Fe}$ values in hematite as high as $+0.75$ to $+1.0\%$ (Johnson *et al.*, 2003) compared with the predicted values for Fe effused from hydrothermal vents ($\pm 0\%$). Consequently, in the case of anoxic Archean oceans, the positive fractionation appears to be consistent with (although not proof of) phototrophic Fe(II) oxidation in the Archean oceans. On the other hand, the similarity in values from anoxygenic phototrophs with those of Fe(III) oxyhydroxides formed by chemical oxidation makes it difficult to distinguish between biotic and abiotic processes (Bullen *et al.*, 2001). Iron isotope fractionation during the oxidation itself is obviously independent of the oxidation mechanism (chemical or biological) because of the very rapid isotopic exchange between aqueous Fe^{2+} and Fe^{3+} . Therefore, isotopic equilibrium will be reached between reduced and oxidized aqueous species.

Plausibility of anoxygenic photoferrotrophy based on Fe and C budget

Ecophysiological studies of these modern strains of anoxygenic Fe(II)-oxidizing phototrophs have been utilized to test their plausibility as a BIF depositional mechanism, with specific regard to carbon and Fe(III) oxyhydroxide productivity. For example, it has been suggested that communities of Fe(II)-oxidizing phototrophs in the early Archean could have generated up to 1.9×10^{13} mol C year⁻¹ throughout all oceans (Kharecha *et al.*, 2005). To put this number into context, Canfield (2005) has estimated that the total net primary productivity in the late Archean–Palaeoproterozoic oceans was 1.8 to 5.6×10^{14} mol C year⁻¹.

By modelling the photosynthetic Fe(II) oxidation rates, it has been suggested that such microorganisms could have accounted for all of the

Fe(III) initially deposited in primary BIF sediment (Konhauser *et al.*, 2002; Kappler & Newman, 2004; Kappler *et al.*, 2005). Due to the pigments they utilize (Bchl and carotenoids), anoxygenic phototrophs absorb light over specific wavelengths (Bchl a 800 to 880 nm, Bchl b 1020 nm, Bchl c ca 750 nm, carotenoids 360 to 517 nm), many of which (>300 nm and <600 nm) would be filtered out in the first few metres of ocean water. By filtering out the wavelengths >650 nm, experimental studies have shown that the oxidation rate of these strains was reduced to 20% of that of the full spectrum, yet that rate still allowed for the oxidation of 0.08 mm Fe(II) per day. Based on this reduction in oxidation efficiency due to light filtering, the average cell density in a planktonic population, assumptions of Archean ocean Fe(II) concentrations (ca 0.5 mM; Holland, 1973; Morris, 1993) and the flux of dissolved Fe(II) from the deep oceans, a conservative estimate for the layer thickness of a planktonic population of anoxygenic Fe(II)-oxidizing phototrophs required to oxidize all dissolved Fe(II) present was 17.6 m (Kappler *et al.*, 2005). With this information, a theoretical amount of Fe(III) oxyhydroxide precipitated by planktonic anoxygenic phototrophs was estimated to be ca 9.0×10^{12} mol Fe yr $^{-1}$ for large basins such the Hamersley in Western Australia (which is ca 10^{11} m 2); this value is consistent with calculations based on maximum BIF deposition rates of 1 mm yr $^{-1}$ where the amount of iron precipitated annually is ca 4.5×10^{12} mol Fe yr $^{-1}$ (Konhauser *et al.*, 2002). Considering the parameters for a model of Fe(II) oxidation rate, light intensity, substrate concentration and cell numbers, even with reduced efficiency in Fe(II) oxidation at conservative values of nutrients and light, anoxygenic phototrophs had the potential to play a significant role in BIF deposition.

ROLE OF CELLULAR CARBON IN POST-DEPOSITIONAL ALTERATION

If a direct biological mechanism was important in the initial process of Fe(II) oxidation in an Archean ocean water column, it is then expected that biomass settled to the sea floor along with the Fe(III) minerals (Konhauser *et al.*, 2011a). Yet, BIFs contain very little organic carbon, meaning that the process was either fully abiotic, or biomass was cycled in the water column or bottom sediment via the

combined metabolic processes of fermentation and chemoheterotrophy (Konhauser *et al.*, 2005). In the case of the latter, any buried organic carbon would be oxidized and, importantly, any cellular biomass would be destroyed, i.e. no microfossils would be preserved in the Fe-rich layers.

If, as today, the organic carbon was oxidized during burial by either diagenesis or metamorphism, the relevant question is which terminal electron acceptor was present at the sea floor during times of BIF deposition, and at what concentrations? Despite the possibility of a surface water oxic zone generated by cyanobacterial activity by 2.7 Ga (see discussion above), deep waters remained anoxic (Canfield, 1998). In the absence of O $_2$, the fermentation products in the bottom waters and/or shallow sediments would have been oxidized via some other form of anaerobic respiratory process (Rothman *et al.*, 2003). In terms of such pathways, the paucity of O $_2$ would have meant minimal nitrate and sulphate availability; the latter being evident from negligible sulphur isotopic fractionations between sulphide and sulphate minerals during the Archean (Strauss, 2003), and the absence of pyrite in BIFs, except in association with inter-layered shaley units (Ewers & Morris, 1981). The supply of MnO $_2$ was probably also not significant because the concentration of Mn(II) released in hydrothermal effluent is up to five times lower than that of iron (Campbell *et al.*, 1988). Furthermore, there are presently no known phototrophic Mn(II)-oxidizing bacteria that would allow the formation of significant amounts of Mn(IV) oxides in the absence of O $_2$. By contrast, however, there was abundant Fe(III) oxyhydroxide deposited as BIFs, and given the presence of partially reduced phases such as magnetite and siderite, a microbial process coupling the oxidation of organic carbon to the reduction of ferric iron producing such reduced iron mineral phases seems very likely (Nealson & Myers, 1990).

Supporting evidence for an ancient Fe(III) reduction pathway comes from the experimental observations that many deeply branching *Archaea* (i.e. some of the oldest purported species) are capable of using H $_2$ to reduce Fe(III) to support chemolithoautotrophic growth (Vargas *et al.*, 1998). These organisms can even use quinone moieties as electron shuttles between solid-phase ferric iron-containing minerals and H $_2$, thereby alleviating the need for direct contact between the cell and mineral

surface (Lovley *et al.*, 2000). Moreover, Fe(III) reduction has been shown to be broadly distributed among several known *Proteobacteria* genera, suggesting that this form of metabolism became widespread over the course of evolution (Barns & Nierzwicki-Bauer, 1997).

Evidence of ancient microbial Fe(III) reduction comes from Fe isotopic ratios in Fe-bearing minerals from Archean sedimentary rocks that closely mimic those observed during modern dissimilatory Fe(III) reduction. For instance, analyses of magnetite-rich rocks from the 2.9 Ga Rietkuil Formation, Witwatersrand Supergroup, South Africa, revealed $\delta^{56}\text{Fe}$ values as low as -2.3‰ (Yamaguchi *et al.*, 2005), whereas siderite in the 2.5 Ga Dales Gorge Member, Brockman Iron Formation, Western Australia, had $\delta^{56}\text{Fe}$ values as low as -2.3‰ (Steinhofel *et al.*, 2010). These negative fractionations are very similar to what might be expected from multiple stages of Fe(III) reduction (Johnson *et al.*, 2004), where each single step during bacterial Fe(III) reduction leads to isotopically light ($\delta^{56}\text{Fe} < 1.2\text{‰}$) aqueous Fe(II) relative to the initial ferric hydroxide substrate (Icopini *et al.*, 2004; Crosby *et al.*, 2007). The importance of Fe(III) reduction in iron fractionation has, however, been challenged by Rouxel *et al.* (2005). These authors suggested that highly variable, but negative, values in pyrite from black shales (0.5 to -3.5‰) between 2.8 Ga and 2.3 Ga is more likely to reflect the initial deposition of Fe(III) oxides (for example, BIFs) that preferentially removed isotopically heavy ^{56}Fe , driving the ocean waters to the negative $\delta^{56}\text{Fe}$ values recorded in pyrite. This interpretation is consistent with the notion that partial biological and abiogenic processes oxidized the dissolved Fe(II) brought into the shallow waters, but is problematic in light of recent studies showing significant isotopic variations for minerals within close proximity, and thus time of deposition (Johnson *et al.*, 2008). As highlighted by Johnson *et al.* (2008), the most plausible way of explaining this variation is if the Fe isotopes reflect diagenetic pathways, and not the Fe composition of the bulk sea water.

The fate of cell biomass (organic carbon) deposited with Fe(III) minerals has not yet been shown experimentally. Theoretically, the relation between Fe(III) and C_{org} in the primary sediment would be stoichiometric if the Fe(II) was oxidized to Fe(III) at the location where biomass was produced, and if the C_{org} and Fe(III) minerals were deposited together, such as in the anoxygenic photoferrotroph model. In this sce-

nario, Fe(III) could be completely re-reduced to Fe(II). This could be the case involving anoxygenic photoferrotrophs. However, from experimental studies it is known that some planktonic cells remain in the water column (Konhauser *et al.*, 2005; Miot *et al.*, 2009; Posth *et al.*, 2010) and only a portion of the cells form aggregates with the Fe(III) hydroxide (Fig. 3), which then settle to the sea floor. The reduction of the Fe(III) mineral and the formation of secondary iron minerals, such as magnetite or siderite, would follow. In such a case, Fe(II) in BIFs can be used as a proxy for deposited C_{org} if the Fe(II) formed by Fe(III) reduction remains in the BIFs.

In contrast to the anoxygenic phototroph model, if the dissolved Fe(II) was oxidized by an abiotic mechanism, either via the UV-photooxidation mechanism or chemical oxidation by cyanobacterially generated O_2 at some distance from the cells (Fig. 2A and B), then the Fe(III) hydroxides produced would be spatially separated from the biomass. This Fe(III) could further react with excess dissolved Fe(II) in the water column forming magnetite [reaction 3], while the cells could be transported away from the shore (Fig. 2B) or be continuously present in the open ocean (Fig. 2C).



As was described for the photoferrotrophs, this scenario could also lead to a partial association of biomass with Fe(III). In this regard, the presence of cyanobacteria or of photoferrotrophs would be difficult to distinguish based on mineralogical or iron isotope studies, but possibly with the identification of metabolism-specific biomarkers.

To estimate the likelihood and plausible impact of the anoxygenic phototroph scenario, Konhauser *et al.* (2005) modelled the Archean marine Fe cycle by making two assumptions. Firstly, it was assumed that the bulk of the Fe(II) component in Fe-rich BIF-type macrobands formed diagenetically through biological Fe(III) reduction, i.e. the magnetite is not primary. Based on a predicted rate of Fe(III) deposition annually (1 mm yr^{-1}), Konhauser *et al.* (2005) then quantified how many electrons were needed to generate that amount of magnetite reported in BIFs (one-third of the ferric oxide minerals; Morris, 1993). Secondly, these authors quantified the amount of photosynthetic Fe(II)-

oxidizer biomass that may have been generated in the photic zone of the water column (based on Kappler *et al.*, 2005) to estimate the amount of Fe recycled prior to burial. The results demonstrated that, under ideal growth conditions, as much as 70% of the biologically formed Fe(III) could have been recycled back into the water column via fermentation and organic carbon oxidation coupled to microbial Fe(III) reduction. Konhauser *et al.* (2005) also suggested that some of the biomass may have been ultimately consumed via methanogenesis, i.e. coupling the oxidation of acetate or H₂ to methane formation. That hypothesis is to some extent corroborated by the analyses of kerogens (extracted from rocks 2.8 Ga and 2.6 Ga) with highly negative $\delta^{13}\text{C}$ signatures (between -40‰ and -60‰) that possibly formed as the result of methanogenic ¹²C-rich gas production, the incorporation of this methane into the biomass of methanotrophic bacteria and inevitably the preservation of ¹²C-enriched organic matter (Hayes, 1983; Eigenbrode & Freeman, 2006).

Recent laboratory experiments by the present authors have focused on the influence of temperature (170°C) and pressure (1.2 kbar) on biogenic Fe(III) minerals associated with biomass by utilizing chemically synthesized ferrihydrite (as a proxy for biogenic ferric hydroxide) and glucose (as a proxy for biomass) mixtures in lieu of biogenic minerals. When chemically synthesized ferrihydrite was treated alone, it was transformed to hematite via dehydration. Mixtures of ferrihydrite and glucose treated under the same temperature and pressure conditions for just a few days produced not only hematite, but also magnetite and siderite. The presence of a cell biomass proxy in low amounts resulted in the production of hematite and small amounts of magnetite and siderite. When the amount of initial biomass proxy was increased, considerable quantities of magnetite and siderite formed while hematite was no longer detectable. These experiments demonstrate that the amount of biomass deposited to the ocean floor in association with the Fe(III) minerals controls the type of Fe minerals preserved in BIFs over geological time frames at elevated temperature and pressure. This shows that the key minerals found in BIF deposits today can be produced, but that there is also a difference in mineral transformation in abiotic and biotic systems (Walker, 1984; Johnson *et al.*, 2003, 2008; Konhauser *et al.*, 2005; Koehler *et al.*, 2013).

BIOLOGICALLY GENERATED SEDIMENTOLOGICAL FEATURES

As described above, many BIFs comprise layers containing magnetite and/or hematite, which alternate on the scale of several millimetres and harbour bands of microcrystalline silica. On average, the microbands range from 0.2 to 1.6 mm in thickness, whereas the total Fe content of the microbands is inversely related to the microband thickness; varying from 30% Fe when the microband is 0.3 mm to 5% when the thickness is 1.5 mm (Trendall & Blockley, 1970). This correlation indicates that the iron content in microbands is constant, whereas variations in the silica content control both the layer thickness and relative iron content (Trendall, 2002). Well-banded iron formations, the typical BIFs, are mostly restricted to Archean and early Palaeoproterozoic sequences (for example, the Transvaal Supergroup, South Africa; Fig. 4A and B). By contrast, large portions of late Palaeoproterozoic iron formations from the Superior and Slave cratons in North America are comprised of sand-sized grains that commonly are cross-bedded and lack the finely laminated textures of BIFs; these shallow-water deposits are generally referred to as granular iron formations (GIFs). These GIFs are typically intercalated with well-laminated BIFs, Fe-rich mudstones, mafic and felsic volcanic rocks, carbonate and sandstone (for example, the Rapitan Formation, Canada; Fig. 4C and D).

The origin of the banding in BIFs remains enigmatic, with two potential end member processes being responsible for their presence. The first assumes that the banding is a primary feature caused by episodic precipitation of the iron layers. The second assumes that the banding was caused by some combination of secondary physical and post-depositional processes. In the case of the former, this can include, for example, BIF precursor sediment being transported to the BIF depositional basin via physical processes, such as density currents (Krapež *et al.*, 2003). In the case of the latter, both allochemical and authigenic BIF sediment would then be subject to chemical and mineralogical transformations arising during burial. The banding thus results when either the Fe(III) was reduced to Fe(II) and the latter diffused out of the sediments, or the silica layers were dissolved and reprecipitated. Although a combination of these processes is likely to have arisen, there still exists a lack of consensus as to which processes were more




Fig. 4. Thin sections of BIF from the Mamatwan Manganese Mine, Hotazel, Northern Cape Province, South Africa, show jasper and magnetite microbanding. This example demonstrates the variation from thin, alternating lens structures (A) at 83.4 m depth to well-developed, uninterrupted and alternating fine layers (B) at 87.5 m depth. In contrast, thin sections from the Rapitan Formation, Canada GIF depict the typical sand-sized grains and discontinuous, uneven layering that suggests stormwave or current influence (C) and (D).

important, and how these processes may have varied from one BIF depositional basin to another.

Primary banding

Based on microbanding in the 2.48 Ga Dales Gorge Member of Western Australia, that was presumably correlated over 100 km (Trendall & Blockley, 1970; Ewers & Morris, 1981), it was proposed that deposition must have been uniform across the basin. Moreover, given the relatively regular spacing between microbands within some larger mesobands, Trendall & Blockley (1970) further proposed that each iron and silica-rich microband couplet resulted from one year of chemical precipitation. The term 'varving' is often used to describe these couplets. It should, however, be pointed out that the iron-rich and silica-rich microbands themselves are often comprised of multiple

internal laminae, which possibly reflects some form of seasonal variations in terms of deposition. For instance, Morris (1993) suggested that the internal complexity may follow four main stages: (i) during the summer, iron was precipitated in the photic zone, producing the iron-rich laminae; (ii) the now depleted photic zone was replenished several times from the subsurface by storm mixing, leading to additional iron-rich laminae; (iii) in the winter, fewer storms meant less Fe replenishment, while cooling led to silica precipitation from sea water that was already saturated with respect to amorphous silica; and (iv) with the return of the summer, iron was once again precipitated.

The link between BIF deposition and seasonality was later advanced by Posth *et al.* (2008) who demonstrated how temperature fluctuations in the surface oceans could link the biotic precipitation of Fe(III) hydroxides by anoxygenic Fe(II)-oxidizing phototrophic bacteria to the

abiotic precipitation of silica. The precipitation of ferric oxyhydroxides by these bacteria is linked directly to their physiological temperature dependence; within a range, these organisms can precipitate more or less ferric oxyhydroxides. For instance, the bacteria grew most profusely in the temperature range of 20 to 25°C, with a decrease in growth rates at lower temperatures and a strict upper temperature limit (Posth *et al.*, 2008). By contrast, silica precipitation is abiotically induced by lowering temperature, and increased solubilization of solid-phase silica at higher temperatures. Furthermore, the extent of temperature fluctuations, for which these processes were demonstrated with model organisms, are within the range experienced in the world oceans today over a yearly cycle, meaning that temperature could have been a unifying trigger for Fe and Si banding.

Other depositional models for the Dales Gorge Member advocate that BIF deposition took place on the ocean basin floor, beyond the slope environment (Krapež *et al.*, 2003). In this regard, it remains unclear what, if any, role microbes may have played in the initial Fe(III) hydroxide precipitation. The model proposes that the Fe(III)-rich sediments accumulated during periods of rising and high sea-level, and presumably were sourced from a submarine hydrothermal system and transported to the BIF depositional basin via deep-sea density currents. The chert layers, by contrast, represent *in situ* silicification of the existing sediment. The chert defines the top of every depositional sequence by sharp eroded contacts with overlying dolostone or mudrock and gradational contacts into underlying BIFs. The chert units also contain multiple shale relicts, 5 to 10 cm thick, which indicates that ambient suspension sediments during sediment starvation (i.e. a drop in sea-level) were not BIFs, and that BIF deposition ceased prior to the end of depositional cycles. Similar features are not only preserved throughout all of the Hamersley Province, but they also have been documented in the Transvaal Province (Beukes & Gutzmer, 2008); it appears to be the common architecture of all deep-water BIFs (Bekker *et al.*, 2010).

Diagenetic banding

A number of studies have suggested that the fine-scale banding in BIFs is a secondary feature derived during burial diagenesis. One of the first

such studies was based largely on the observation of compressed lateral terminations of chert pods (Trendall & Blockley, 1970). These authors demonstrated that the compressed areas are enriched in iron minerals at the expense mainly of silica and, using this evidence, concluded that all high iron bands, podded or not, represented compression of the primary BIF sediment with vertical escape of silica. The remobilized silica then reprecipitated evenly into overlying pods that are rich in silica. Along similar lines, Simonson (1987) proposed that the chert pods were concretions that formed early via local cementation, followed by differential compaction of the surrounding, less indurated sediments during burial.

More recently, and based instead on facies and sequence-stratigraphic analyses, Krapež *et al.* (2003) and Pickard *et al.* (2004) concluded that all chert in BIFs is diagenetic in origin. Moreover, these authors concluded that chert mesobands are siliceous equivalents of modern-day sea floor hardgrounds, in which silica replaced precursor sediment at or below the sediment–water interface. Three-dimensional and micro-scale lenticularity of chert and relicts of precursor sediment within lamina sets and discontinuous bands, as well as erosion surfaces on bedded cherts, show that chert has a replacement origin and formed during early diagenesis, prior to compaction. As discussed above, the iron layers, by contrast, represent re-suspended Fe(III) hydroxide-rich sediments that were carried into the depositional basin via density currents and then subject to post-depositional processes (Krapež *et al.*, 2003).

Other studies have put forward the possibility that it was the iron that was selectively mobilized during diagenesis. Based on the earlier discovery that some anaerobic heterotrophs could grow by coupling the oxidation of organic matter to the reduction of Fe(III) hydroxides (Lovley & Phillips, 1988), Nealson & Myers (1990) subsequently proposed that such bacteria could provide a mechanism by which iron-poor layers could be generated during diagenesis; the iron-rich layers represent sediment buried in which molar Fe : C ratios exceeded unity and all the biomass was oxidized. Coupling the reduction of Fe(III) minerals to the oxidation of organic matter not only explains the low content of organic carbon in the BIFs via the consumption of carbon (<0.5%; Gole & Klein, 1981), but also explains the abundance of light carbon isotopic signatures associated with the interlayered

carbonate minerals (Perry *et al.*, 1973; Walker, 1984; Baur *et al.*, 1985).

CONCLUSIONS

Understanding the mechanism(s) underpinning banded iron formation (BIF) deposition would offer significant insights into some of the biogeochemical processes that took place on the early Earth. The first suggestions of a microbial role were postulated over 40 years ago (Cloud, 1965). At present, there is still indirect evidence of a microbial mechanism driven by either anoxygenic Fe(II)-oxidizing photosynthesis or cyanobacterially driven O₂ production. Moreover, the complexity of BIF deposition, on temporal and spatial scales, makes it possible that multiple mechanisms aided BIF deposition, depending on the ocean and atmospheric redox state, as well as on the local geochemical and microbiological conditions. One could imagine a scenario with anoxygenic phototrophs being mainly responsible for the deposition of the BIFs in low O₂ oceans prior to the Great Oxidation Event (GOE). After the emergence of cyanobacteria, it has been suggested that O₂ producing cyanobacteria could have helped drive the abiotic oxidation of Fe(II), whereas anoxygenic phototrophs carried out Fe(II) oxidation in an anoxic niche (Kappler *et al.*, 2005). In order to understand BIFs and the biogeosphere in which they were deposited, as well as to be able to interpret them as an archive of Earth history, it is necessary to take a holistic approach – integrating new methods for biomarker and stable isotope analyses, and also keeping this in context with mineralogical examinations and ecophysiological studies with modern analogue micro-organisms.

ACKNOWLEDGEMENTS

We thank Nicolas Beukes for his advice on South African BIFs, as well as his guidance in sample collection. Gert van der Linde of the Hotazel Manganese Mines, Hotazel, South Africa, provided BIF samples. We also thank Claus Burkhardt and Sebastian Schädler for the preparation of SEM images, which was carried out at the Natural and Medical Sciences Institute at the University of Tuebingen (NMI). This work was supported by an Emmy-Noether fellowship and a research grant from the German Research Foundation (DFG) made to AK (KA

1736/2-1, 2-2 and 4-1), a research grant from the German Research Foundation (DFG) made to N.P. (PO-1624/1-1, 2-1), and a Natural Sciences and Engineering Research Council award to KOK. We thank Clark Johnson, Phil Fralick, Nora Noffke and an anonymous reviewer for their helpful comments, which greatly improved the quality of the manuscript.

REFERENCES

- Altermann, W. and Kazmierczak, J. (2003) Archean microfossils: a reappraisal of early life on Earth. *Res. Microbiol.*, **154**, 611–617.
- Altermann, W. and Schopf, J.W. (1995) Microfossils from the Neoproterozoic Campbell Group, Griqualand West Sequence of the Transvaal Supergroup, and their paleoenvironmental and evolutionary implications. *Precambrian Res.*, **75**, 65–90.
- Anbar, A.D., Duan, Y., Lyons, T.W., Arnold, G.L., Kendall, B., Creaser, R.A., Kaufman, A.J., Gordon, G.W., Scott, C., Garvin, J. and Buick, R. (2007) A whiff of oxygen before the great oxidation event? *Science*, **317**, 1903–1906.
- Balci, N., Bullen, T.D., Witte-Lien, K., Shanks, W.C., Motelica, M. and Mandernack, K.W. (2006) Iron isotope fractionation during microbially stimulated Fe(II) oxidation and Fe(III) precipitation. *Geochim. Cosmochim. Acta*, **70**, 622–639.
- Barns, S.M. and Nierzwicki-Bauer, S.A. (1997) Microbial diversity in ocean, surface and subsurface environments. In: *Geomicrobiology: Interactions Between Microbes and Minerals* (Eds J.F. Banfield and K.H. Nealson), pp. 35–79. Mineralogical Society of America, Washington, DC.
- Bau, M. and Möller, P. (1993) Rare earth element systematics of the chemically precipitated component in Early Precambrian iron-formations and the evolution of the terrestrial atmosphere-hydrosphere-lithosphere system. *Geochim. Cosmochim. Acta*, **57**, 2239–2249.
- Baur, M.E., Hayes, J.M., Studley, S.A. and Walter, M.R. (1985) Millimeter-scale variations of stable isotope abundances in carbonates from banded iron formations in the Hamersley Group of Western Australia. *Econ. Geol.*, **80**, 270–282.
- Bekker, A., Holland, H.D., Wang, P.-L., Rumble, D., III, Stein, H.J., Hannah, J.L., Coetzee, L.L. and Beukes, N.J. (2004) Dating the rise of atmospheric oxygen. *Nature*, **427**, 117–120.
- Bekker, A., Slack, J.F., Planavsky, N., Krapež, B., Hofmann, A., Konhauser, K.O. and Rouxel, O.J. (2010) Iron formation: a sedimentary product of the complex interplay among mantle, tectonic, and biospheric processes. *Econ. Geol.*, **105**, 467–508.
- Beukes, N.J. (1973) Precambrian iron-formations of Southern Africa. *Econ. Geol.*, **68**, 960–1004.
- Beukes, N.J. (1984) Sedimentology of the Kuruman and Griquatown Iron-Formations, Transvaal Supergroup, Griqualand West, South Africa. *Precambrian Res.*, **24**(1), 47–84.
- Beukes, N.J. and Gutzmer, J. (2008) Origin and paleoenvironmental significance of major iron formations at the Archean-Paleoproterozoic boundary. *Rev. Econ. Geol.*, **15**, 5–47.

- Bosak, T., Greene, S.E. and Newman, D.K.** (2007) A possible role for anoxygenic photosynthetic microbes in the formation of ancient stromatolites. *Geobiology*, **5**, 119–126.
- Bosak, T., Liang, B., Sim, M.S. and Petroff, A.P.** (2009) Morphological record of oxygenic photosynthesis in conical stromatolites. *Proc. Natl Acad. Sci. USA*, **106**(27), 10939–10943.
- Brasier, M.D., Green, O.R., Jephcoat, A.P., Kleppe, A.K., Van Kranendonk, M.J., Lindsay, J.F., Steele, A. and Grassineau, N.V.** (2002) Questioning the evidence for Earth's oldest fossils. *Nature*, **416**, 76–81.
- Brasier, M.D., Green, P.R., Lindsay, J.F., McLoughlin, N., Steele, A. and Stoakes, C.** (2005) Critical testing of Earth's oldest putative fossil assemblage from the ~3.5 Apex chert, Chinaman Creek, Western Australia. *Precambrian Res.*, **140**, 55–105.
- Braterman, P.S., Cairns-Smith, A.G. and Sloper, R.W.** (1983) Photo-oxidation of hydrated Fe²⁺ - significance for banded iron formations. *Nature*, **303**, 163–164.
- Brocks, J.J.** (2005) Biomarker evidence for green and purple sulphur bacteria in a stratified Palaeoproterozoic sea. *Nature*, **437**, 866–870.
- Brocks, J.J., Logan, G.A., Buick, R. and Summons, R.E.** (1999) Archean molecular fossils and the early rise of eukaryotes. *Science*, **285**, 1033–1036.
- Buick, R.** (1988) Carbonaceous filaments from North Pole, West Australia: are they fossil bacteria in Archean stromatolites? A reply. *Precambrian Res.*, **39**, 311–317.
- Buick, R.** (1992) The antiquity of oxygenic photosynthesis: evidence for stromatolites in sulphate-deficient Archean lakes. *Science*, **255**, 74–77.
- Buick, R.** (2008) When did oxygenic photosynthesis evolve? *Philos. Trans. R. Soc. Lond. B: Biol. Sci.*, **363**, 2731–2743.
- Bullen, T.D., White, A.F., Childs, C.W., Vivit, D.V. and Schulz, M.J.** (2001) Demonstration of significant abiotic iron isotope fractionation in nature. *Geology*, **29**(8), 699–702.
- Burns, B.P., Goh, F., Allen, M. and Neilan, B.A.** (2004) Microbial diversity of extant stromatolites in the hypersaline marine environment of Shark Bay, Australia. *Environ. Microbiol.*, **6**, 1096–1101.
- Burns, B.P., Anitori, R., Butterworth, P., Henneberger, R., Goh, F., Allen, M.A., Ibañez-Peral, R., Bergquist, P.L., Walter, M.R. and Neilan, B.A.** (2009) Modern analogues and the early history of microbial life. *Precambrian Res.*, **173**, 10–18.
- Cairns-Smith, A.G.** (1978) Precambrian solution photochemistry, inverse segregation, and banded iron formations. *Nature*, **276**, 807–808.
- Campbell, A.C., Palmer, M.R., Klinkhammer, G.P., Bowers, T.S., Edmond, J.M., Lawrence, J.R., Casey, J.F., Thompson, G., Humphris, S., Rona, P. and Karson, J.A.** (1988) Chemistry of hot springs on the Mid-Atlantic Ridge. *Nature*, **335**, 514–519.
- Canfield, D.E.** (1998) A new model for Proterozoic ocean chemistry. *Nature*, **396**, 450–453.
- Canfield, D.E.** (2005) The early history of atmospheric oxygen: homage to Robert M. Garrels. *Annu. Rev. Earth Planet. Sci.*, **33**, 1–36.
- Canfield, D.E., Habicht, K.S. and Thamdrup, B.** (2000) The Archean sulfur cycle and the early history of atmospheric oxygen. *Science*, **288**, 658–661.
- Cloud, P.** (1965) Significance of the Gunflint (Precambrian) microflora. *Science*, **148**, 27–35.
- Cloud, P.** (1969) Atmospheric and hydrospheric evolution on the primitive Earth. *Science*, **160**, 729–736.
- Cloud, P.** (1973) Paleocological significance of the banded iron-formation. *Econ. Geol.*, **68**, 1135–1143.
- Croal, L.R., Johnson, C.M., Beard, B.L. and Newman, D.K.** (2004) Iron isotope fractionation by Fe(II)-oxidizing photoautotrophic bacteria. *Geochim. Cosmochim. Acta*, **68**, 1227–1242.
- Croal, L.R., Jiao, Y., Kappler, A. and Newman, D.K.** (2009) Phototrophic Fe(II) oxidation in an atmosphere of H₂: implications for Archean banded iron formations. *Geobiology*, **7**(1), 21–24.
- Crosby, H.A., Roden, E.E., Johnson, C.M. and Beard, B.L.** (2007) The mechanisms of iron isotope fractionation produced during dissimilatory Fe(III) reduction by *Shewanella putrefaciens* and *Geobacter sulfurreducens*. *Geobiology*, **5**, 169–189.
- Dimroth, E. and Chauvel, J.J.** (1973) Petrography of the Sokoman iron formation in part of the central Labrador trough, Quebec, Canada. *Geol. Soc. Am. Bull.*, **84**, 111–134.
- Ehrenreich, A. and Widdel, F.** (1994) Anaerobic oxidation of ferrous iron by purple bacteria, a new-type of phototrophic metabolism. *Appl. Environ. Microbiol.*, **60**, 4517–4526.
- Eickhoff, M., Birgel, D., Talbot, H.M., Peckmann, J. and Kappler, A.** (2013) Oxidation of Fe(II) leads to increased C-2 methylation of pentacyclic triterpenoids in the anoxygenic phototrophic bacterium *Rhodospseudomonas palustris* strain TIE-1. *Geobiology*, **11**, 268–278.
- Eigenbrode, J.L. and Freeman, K.H.** (2006) Late Archean rise of aerobic microbial ecosystems. *Proc. Natl Acad. Sci. USA*, **103**(43), 15759–15764.
- Eigenbrode, J.L., Freeman, K.H. and Summons, R.E.** (2008) Methylhopane biomarker hydrocarbons in Hamersley Province sediments provide evidence for Neoproterozoic aerobicity. *Earth Planet. Sci. Lett.*, **273**, 323–331.
- Ewers, W.E. and Morris, R.C.** (1981) Studies of the Dales Gorge Member of the Brockman iron formation, Western Australia. *Econ. Geol.*, **76**, 1929–1953.
- Farquhar, J., Bao, H. and Thiemens, M.** (2000) Atmospheric influence of Earth's earliest sulfur cycle. *Science*, **289**, 756–758.
- Farquhar, J.F., Zerkle, A.K. and Bekker, A.** (2011) Geological constraints on the origin of oxygenic photosynthesis. *Photosynth. Res.*, **107**, 11–36.
- Fischer, W.W. and Pearson, A.** (2007) Hypotheses for the origin and early evolution of triterpenoid cyclases. *Geobiology*, **5**, 19–34.
- Fischer, W.W., Summons, R.E. and Pearson, A.** (2005) Targeted genomic detection of biosynthetic pathways: anaerobic production of hopanoid biomarkers by a common sedimentary microbe. *Geobiology*, **3**, 33–40.
- Fischer, W.W., Schroeder, S., Lacassie, J.P., Beukes, N.J., Goldberg, T., Strauss, H., Horstmann, U.E., Schrag, D.P. and Knoll, A.H.** (2009) Isotopic constraints on the late Archean carbon cycle from the Transvaal Supergroup along the western margin of the Kaapvaal Craton, South Africa. *Precambrian Res.*, **169**, 15–27.
- Fralick, P. and Pufahl, P.K.** (2006) Iron formation in neoproterozoic deltaic successions and the microbially mediated deposition of transgressive systems tracts. *J. Sed. Res.*, **76**, 1057–1066.

- Frei, R., Gaucher, C., Poulton, S.W. and Canfield, D.E. (2009) Fluctuations in Precambrian atmospheric oxygenation recorded by chromium isotopes. *Nature*, **461**, 250–254.
- Garcia-Ruiz, J.M., Hyde, S.T., Carnerup, A.M., Christy, A.G., Van Kranendonk, M.J. and Welham, N.J. (2003) Self-assembled silica-carbonate structures and detection of ancient microfossils. *Science*, **302**, 1194–1197.
- Garrels, R.M., Perry, E.A., Jr and MacKenzie, F.T. (1973) Genesis of Precambrian iron-formations and the development of atmospheric oxygen. *Econ. Geol.*, **68**, 1173–1179.
- Godfrey, L.V. and Falkowski, P.G. (2009) The cycling and redox state of nitrogen in the Archaean ocean. *Nature Geosci.*, **2**, 725–729.
- Gole, M.J. and Klein, C. (1981) Banded Iron Formations through much of Precambrian time. *J. Geol.*, **89**, 169–183.
- Grotzinger, J.P. and Knoll, A.H. (1999) Stromatolites in Precambrian carbonates: evolutionary mileposts or environmental dipsticks? *Annu. Rev. Earth Planet. Sci.*, **27**, 313–358.
- Grotzinger, J.P. and Rothman, D.H. (1996) An abiotic model for stromatolite morphogenesis. *Nature*, **383**, 423–425.
- Hamade, T., Konhauser, K.O., Raiswell, R., Goldsmith, S. and Morris, R.C. (2003) Using Ge/Si ratios to decouple iron and silica fluxes in Precambrian banded iron formations. *Geology*, **31**, 35–38.
- Han, T.-M. (1966) Textural relations of hematite and magnetite in some Precambrian metamorphosed oxide iron-formations. *Econ. Geol.*, **61**, 1306–1310.
- Han, T.-M. (1978) Microstructures of magnetite as guides to its origin in some Precambrian iron-formations. *Fortschr. Mineral.*, **56**(1), 105–142.
- Hartman, H. (1984) The evolution of photosynthesis and microbial mats: a speculation on banded iron formations. In: *Microbial Mats: Stromatolites* (Eds Y. Cohen, R.W. Castenholz and H.O. Halvorson), pp. 451–453. Alan Liss, New York.
- Härtner, T., Straub, K.L. and Kannenberg, E. (2005) Occurrence of hopanoid lipids in anaerobic *Geobacter* species. *FEMS Microbiol. Lett.*, **243**, 59–64.
- Hayes, J.M. (1983) Geochemical evidence bearing on the origin of aerobiosis, a speculative hypothesis. In: *Earth's Earliest Biosphere, Its Origins and Evolution* (Ed. J.W. Schopf), pp. 291–301. Princeton University Press, Princeton, NJ.
- Hegler, F., Posth, N.R., Jiang, J. and Kappler, A. (2008) Physiology of phototrophic iron(II)-oxidizing bacteria: implications for modern and ancient environments. *FEMS Microbiol. Ecol.*, **66**(2), 250–260.
- Heising, S. and Schink, B. (1998) Phototrophic oxidation of ferrous iron by a *Rhodomicrobium vannielii* strain. *Microbiology*, **144**, 2263–2269.
- Heising, S., Richter, L., Ludwig, W. and Schink, B. (1999) *Chlorobium ferrooxidans* sp. nov., a phototrophic green sulfur bacterium that oxidizes ferrous iron in coculture with a *Geospirillum* sp. strain. *Arch. Microbiol.*, **172**, 116–124.
- Hoefs, J. (1997) *Stable Isotope Geochemistry*, 4th edn. Springer Verlag, Berlin Heidelberg, 201 pp.
- Holland, H.D. (1973) The oceans: a possible source of iron in iron-formations. *Econ. Geol.*, **68**, 1169–1172.
- Holm, N.G. (1989) The ¹³C/¹²C ratios of siderite and organic matter of a modern metalliferous hydrothermal sediment and their implications for banded iron formations. *Chem. Geol.*, **77**, 41–45.
- Icopini, G.A., Anbar, A.D., Ruebush, S.R., Tien, M. and Brantley, S.L. (2004) Iron isotope fractionation during microbial reduction of iron. *Geology*, **32**(3), 205–208.
- Isley, A.E. (1995) Hydrothermal plumes and the delivery of iron to banded iron formation. *J. Geol.*, **103**, 169–185.
- Isley, A.E. and Abbot, D.H. (1999) Plume-related mafic volcanism and the deposition of banded iron formation. *J. Geophys. Res.*, **104**, 15461–15477.
- Jacobsen, S.B. and Pimentel-Klose, M.R. (1988) A Nd isotopic study of the Hamersley and Michipicoten banded iron formations: the source of REE and Fe in Archaean oceans. *Earth Planet. Sci. Lett.*, **87**, 29–44.
- James, H.L. (1954) Sedimentary facies of iron-formation. *Econ. Geol.*, **49**(3), 236–294.
- Jiao, Y., Kappler, A., Croal, L.R. and Newman, D.K. (2005) Isolation and characterization of a genetically traceable photoautotrophic Fe(II)-oxidizing bacterium, *Rhodospseudomonas palustris* strain Tie-1. *Appl. Environ. Microbiol.*, **71**, 4487–4496.
- Johnson, C.M. and Beard, B.L. (2005) Biogeochemical cycling of iron isotopes. *Science*, **309**, 1025–1027.
- Johnson, C.M., Beard, B.L., Beukes, N.J., Klein, C. and O'Leary, J.M. (2003) Ancient geochemical cycling in the Earth as inferred from Fe isotope studies of banded iron formations from the Transvaal craton. *Contrib. Mineral. Petrol.*, **144**, 523–547.
- Johnson, C.M., Beard, B.L., Roden, E.E., Newman, D.K. and Neilson, K.H. (2004) Isotopic constraints on biogeochemical cycling of Fe. *Rev. Mineral. Geochem.*, **55**, 359–408.
- Johnson, C.M., Roden, E.E., Welch, S.A. and Beard, B.L. (2005) Experimental constraints on Fe isotope fractionation during magnetite and Fe carbonate formation coupled to dissimilatory hydrous ferric oxide reduction. *Geochim. Cosmochim. Acta*, **69**, 963–993.
- Johnson, C.M., Beard, B.L., Klein, C., Beukes, N.J. and Roden, E.E. (2008) Iron isotopes constrain biologic and abiologic processes in banded iron formation genesis. *Geochim. Cosmochim. Acta*, **72**, 151–169.
- Kappler, A. and Newman, D.K. (2004) Formation of Fe(III) minerals by Fe(II) oxidizing photoautotrophic bacteria. *Geochim. Cosmochim. Acta*, **68**, 1217–1226.
- Kappler, A., Pasquero, C., Konhauser, K.O. and Newman, D.K. (2005) Deposition of banded iron formations by anoxygenic phototrophic Fe(II)-oxidizing bacteria. *Geology*, **33**, 865–868.
- Kappler, A., Johnson, C.M., Crosby, H.A., Beard, B.L. and Newman, D.K. (2010) Evidence for equilibrium iron isotope fractionation by nitrate-reducing iron(II)-oxidizing bacteria. *Geochim. Cosmochim. Acta*, **74**, 2826–2842.
- Kaufman, A.J., Johnston, D.T., Farquhar, J., Masterson, A.L., Lyons, T.W., Bates, S., Anbar, A.D., Arnold, G.L., Garvin, J. and Buick, R. (2007) Late Archaean biospheric oxygenation and atmospheric evolution. *Science*, **317**, 1900–1903.
- Kendall, B., Reinhard, C.T., Lyons, T.W., Kaufman, A.J., Poulton, S.W. and Anbar, A.D. (2010) Pervasive oxygenation along late Archaean ocean margins. *Nature Geosci.*, **3**, 647–652.
- Kharecha, P., Kasting, J. and Siefert, J. (2005) A coupled atmosphere-ecosystem model of the early Archaean Earth. *Geobiology*, **3**, 53–76.
- Klein, C. (2005) Some Precambrian Banded Iron Formations (BIFs) from around the world: their age, geologic setting,

- mineralogy, metamorphism, geochemistry, and origin. *Am. Mineral.*, **90**, 1473–1499.
- Klein, C. and Beukes, N.J.** (1992) Time Distribution, Stratigraphy, and Sedimentologic Setting, and Geochemistry of Precambrian Iron-Formations. In: *The Proterozoic Biosphere: A Multidisciplinary Study* (Eds J.W. Schop and C. Klein), pp. 139–146. Press syndicate of the University of Cambridge, Cambridge.
- Koehler, I., Papineau, D., Konhauser, K.O. and Kappler, A.** (2013) Biological carbon precursor to diagenetic siderite spherulites in banded iron formations. *Nature Commun.*, doi:10.1038/ncomms2770.
- Konhauser, K.O.** (2007) *Introduction to Geomicrobiology*. Blackwell, Oxford, 425 pp.
- Konhauser, K., Hamade, T., Raiswell, R., Morris, R.C., Ferris, F.G., Southam, G. and Canfield, D.E.** (2002) Could bacteria have formed the Precambrian banded iron formations? *Geology*, **30**, 1079–1082.
- Konhauser, K., Newman, D.K. and Kappler, A.** (2005) The potential significance of microbial Fe(III) reduction during deposition of Precambrian banded iron formations. *Geobiology*, **3**, 167–177.
- Konhauser, K.O., Amskold, L., Lalonde, S.V., Posth, N.R., Kappler, A. and Anbar, A.** (2007a) Decoupling photochemical Fe(II) oxidation from shallow-water deposition. *Earth Planet. Sci. Lett.*, **258**, 87–100.
- Konhauser, K.O., Lalonde, S., Amskold, L. and Holland, H.D.** (2007b) Was there really an Archean phosphate crisis? *Science*, **315**, 1234.
- Konhauser, K.O., Kappler, A. and Roden, E.E.** (2011a) Iron in microbial metabolism. *Elements*, **7**, 89–93.
- Konhauser, K.O., Lalonde, S.V., Planavsky, N., Pecoits, E., Lyons, T., Mojzsis, S., Rouxel, O.J., Barley, M., Rosiere, C., Fralick, P.W., Kump, L.R. and Bekker, A.** (2011b) Chromium enrichment in iron formations record Earth's first acid rock drainage during the Great Oxidation Event. *Nature*, **478**, 369–373.
- Krapež, B., Barley, M.E. and Pickard, A.L.** (2003) Hydrothermal and resedimented origins of the precursor sediments to banded iron formation: sedimentological evidence from the Early Paleoproterozoic Brockman Supersequence of Western Australia. *Sedimentology*, **50**, 979–1011.
- Kump, L. and Barley, M.E.** (2007) Increased subaerial volcanism and the rise of atmospheric oxygen 2.5 billion years ago. *Nature*, **448**, 1033–1036.
- Kump, L.R. and Holland, H.D.** (1992) Iron in Precambrian rocks: implications for the global oxygen budget of the ancient Earth. *Geochim. Cosmochim. Acta*, **56**, 3217–3223.
- Li, J.L., Konhauser, K.O., Cole, D.R. and Phelps, T.J.** (2011) Mineral ecophysiological data provide growing evidence for microbial activity in banded-iron formations. *Geology*, **39**, 707–710.
- Lovley, D.R. and Phillips, E.J.P.** (1988) Novel mode of microbial energy metabolism: organic carbon oxidation coupled to dissimilatory reduction of iron or manganese. *Appl. Environ. Microbiol.*, **54**, 1472–1480.
- Lovley, D.R., Kashefi, K., Vargas, M., Tor, J.M. and Blunt-Harris, E.L.** (2000) Reduction of humic substances and Fe (III) by hyperthermophilic microorganisms. *Chem. Geol.*, **169**, 289–298.
- Maliva, R.G., Knoll, A.H. and Simonson, B.M.** (2005) Secular change in the Precambrian silica cycle: insights from chert petrology. *Geol. Soc. Am. Bull.*, **117**, 835–845.
- Manske, A.K., Glaeser, J., Kuypers, M.A.M. and Overmann, J.** (2005) Physiology and phylogeny of green sulfur bacteria forming a monospecific phototrophic assemblage at a depth of 100 meters in the Black Sea. *Appl. Environ. Microbiol.*, **71**, 8049–8060.
- Marshall, C.P., Emry, J.R. and Olcott Marshall, A.** (2011) Haematite pseudomicrofossils present in the 3.5-billion-year-old Apex chert. *Nature Geosci.*, **4**, 240–243.
- McLoughlin, N., Wilson, L.A. and Brasier, M.D.** (2008) Growth of synthetic stromatolites and wrinkle structures in the absence of microbes – implications for the early fossil record. *Geobiology*, **6**, 95–105.
- Miot, J., Benzerara, K., Obst, M., Kappler, A., Hegler, F., Schädler, S., Bouchez, C., Guyot, F. and Morin, G.** (2009) Extracellular iron biomineralization by photoautotrophic iron-oxidizing bacteria. *Appl. Environ. Microbiol.*, **75**, 5586–5591.
- Mojzsis, S.J., Coath, C.D., Greenwood, J.P., McKeegan, K.D. and Harrison, T.M.** (2003) Mass-independent isotope effects in Archean (2.5 to 3.8 Ga) sedimentary sulfides determined by ion microprobe analysis. *Geochim. Cosmochim. Acta*, **67**, 1635–1658.
- Morris, R.C.** (1993) Genetic modelling for banded iron-formation of the Hamersley Group, Pilbara Craton, Western Australia. *Precambrian Res.*, **60**, 243–286.
- Morris, R.C. and Horwitz, R.C.** (1983) The origin of the iron-formation-rich Hamersley Group of Western Australia – Deposition on a platform. *Precambrian Res.*, **21**, 273–297.
- Nealson, K. H. and Myers, C. R.** (1990) Iron reduction by bacteria: a potential role in the genesis of banded iron formations. *Am. J. Sci.*, **290-A**, 35–45.
- Noffke, N., Beukes, N., Bower, D., Hazen, R.M. and Swift, D.J.P.** (2008) An actualistic perspective into Archean worlds – (cyano)bacterially induced sedimentary structures in the siliciclastic Nhlazatse Section, 2.9 Ga Pongola Supergroup, South Africa. *Geobiology*, **6**, 5–20.
- Olson, J.M. and Blankenship, R.E.** (2004) Thinking about the evolution of photosynthesis. *Photosynth. Res.*, **80**, 373–386.
- Overmann, J., Cypionka, H. and Pfennig, N.** (1992) An extremely low light adapted phototrophic sulfur bacterium from the Black Sea. *Limnol. Oceanogr.*, **37**, 150–155.
- Papineau, D., Walker, J.J., Mojzsis, S.J. and Pace, N.R.** (2005) Composition and structure of Microbial communities from stromatolites of Hamelin pool in Shark Bay, Western Australia. *Appl. Environ. Microbiol.*, **71**(8), 4822–4832.
- Pasteris, J.D. and Wopenka, B.** (2003) Necessary, but not sufficient Raman identification of disordered carbon as a signature of ancient life. *Astrobiology*, **3**, 727–738.
- Pavlov, A.A. and Kasting, J.F.** (2002) Mass-independent fractionation of sulfur isotopes in Archean sediments: strong evidence for an anoxic Archean atmosphere. *Astrobiology*, **2**, 27–41.
- Perry, E.C., Tan, F.C. and Morey, G.B.** (1973) Geology and stable isotope geochemistry of the Biwabik Iron Formation, northern Minnesota. *Econ. Geol.*, **68**, 1110–1125.
- Phoenix, V.R., Konhauser, K.O., Adams, D.G. and Bottrell, S.H.** (2001) Role of biomineralization as an ultraviolet shield: implications for Archean life. *Geology*, **29**, 823–826.
- Pickard, A.L., Barley, M.E. and Krapež, B.** (2004) Deep-marine depositional setting of banded iron formation: sedimentological evidence from interbedded clastic sedimentary rocks in the early Palaeoproterozoic Dales Gorge Member of Western Australia. *Sed. Geol.*, **170**, 37–62.

- Pierson, B.K. (1994) The emergence, diversification, and role of photosynthetic eubacteria. In: *Early Life on Earth* (Ed. S. Bengtson), pp. 161–180. Nobel Symposium No. 84 held at Björkborn, Sweden, 16–21 May 1992.
- Planavsky, N., Rouxel, O., Bekker, A., Shapero, R., Fralick, P. and Knudsen, A. (2009) Iron-oxidizing microbial ecosystems thrived in late Paleoproterozoic redox-stratified oceans. *Earth Planet. Sci. Lett.*, **286**(1–2), 230–242.
- Posth, N.R., Hegler, F., Konhauser, K.O. and Kappler, A. (2008) Alternating Si and Fe deposition caused by temperature fluctuations in Precambrian oceans. *Nature Geosci.*, **1**(10), 703–708.
- Posth, N.R., Huelin, S., Konhauser, K.O. and Kappler, A. (2010) Size, density and composition of cell-mineral aggregates formed during anoxygenic phototrophic Fe(II) oxidation: impact on modern and ancient environments. *Geochim. Cosmochim. Acta*, **74**, 3476–3493.
- Rashby, S.E., Sessions, A.L., Summons, R.E. and Newman, D.K. (2007) Biosynthesis of 2-methylbacteriohopanepolyols by an anoxygenic phototroph. *Proc. Natl Acad. Sci. USA*, **104**(38), 15099–15104.
- Rasmussen, B. and Buick, R. (1999) Redox state of the Archean atmosphere: evidence from detrital heavy metals in ca. 3250–2750 Ma sandstones from the Pilbara Craton, Australia. *Geology*, **27**, 115–118.
- Rasmussen, B., Fletcher, I.R., Brocks, J.J. and Kilburn, M.R. (2008) Reassessing the first appearance of eukaryotes and cyanobacteria. *Nature*, **455**, 1101–1104.
- Raymond, J. and Blankenship, R.E. (2004) Biosynthetic pathways, gene replacement and the antiquity of life. *Geobiology*, **2**, 199–203.
- Reinhard, C.T., Raiswell, R., Scott, C., Anbar, A.D. and Lyons, T.W. (2009) A Late Archean sulfidic sea stimulated by early oxidative weathering of the continents. *Science*, **326**, 713–716.
- Rothman, D.H., Hayes, J.M. and Summons, R.E. (2003) Dynamics of the Neoproterozoic carbon cycle. *Proc. Natl Acad. Sci. USA*, **100**, 8124–8129.
- Rouxel, O.J., Bekker, A. and Edwards, K.J. (2005) Iron isotope constraints on the Archean and Paleoproterozoic. *Science*, **307**, 1088–1091.
- Rye, R. and Holland, H.D. (1998) Paleosols and the evolution of atmospheric oxygen: a critical review. *Am. J. Sci.*, **298**, 621–672.
- Schaedler, S., Burkhardt, C., Hegler, F., Straub, K.L., Miot, J., Benzerara, K. and Kappler, A. (2009) Formation of cell-iron-mineral aggregates by phototrophic and nitrate-reducing anaerobic Fe(II)-oxidizing bacteria. *Geomicrob. J.*, **26**, 93–103.
- Schidlowski, M. (2000) Carbon isotopes and microbial sediments. In: *Microbial Sediments* (Eds R. Riding and S.M. Awramik), pp. 84–95. Springer Verlag, Berlin.
- Schopf, J.W. (1993) Microfossils of the Early Archean Apex chert: new evidence of the antiquity of life. *Science*, **260**, 640–646.
- Schopf, J.W. and Kudryavtsev, A.B. (2009) Confocal laser scanning microscopy and Raman imagery of ancient microscopic fossils. *Precambrian Res.*, **173**, 39–49.
- Schopf, J.W. and Packer, B.M. (1987) Early Archean (3.3-billion to 3.5-billion-year-old) microfossils from Warrawoona Group, Australia. *Science*, **237**, 70–73.
- Schopf, J.W., Kudryavtsev, A.B., Agresti, D.G., Wdowiak, T.J. and Czaja, A.D. (2002) Laser-Raman imagery of Earth's earliest fossils. *Nature*, **416**, 73–76.
- Schopf, J.W., Kudryavtsev, A.B., Czaja, A.D. and Tripathi, A.B. (2007) Evidence of Archean life: stromatolites and microfossils. *Precambrian Res.*, **158**, 141–155.
- Siever, R. (1992) The silica cycle in the Precambrian. *Geochim. Cosmochim. Acta*, **56**, 3265–3272.
- Simonson, B.M. (1987) Early silica cementation and subsequent diagenesis in arenites from four early Proterozoic iron formations of North America. *J. Sed. Petrol.*, **57**, 494–511.
- Steinhofel, G., von Blackenburg, F., Horn, I., Konhauser, K.O., Beukes, N. and Gutzmer, J. (2010) Deciphering formation processes of banded iron formations from the Transvaal and the Hamersley Sequence by combined Si and Fe isotope analysis using UV femtosecond laser ablation. *Geochim. Cosmochim. Acta*, **74**, 2677–2696.
- Straton, S., Amskold, L., Anbar, A. and Konhauser, K.O. (2006) *Iron Isotope Fractionation During Fe(II) Photo-Oxidation*. Astrobiology Science Conference 4, Washington, DC.
- Straub, K.L., Rainey, F.R. and Widdel, F. (1999) *Rhodovulum iodolum* sp. nov. and *Rhodovulum robiginosum* sp. nov., two new marine phototrophic ferrous-iron-oxidizing purple bacteria. *Int. J. Syst. Bacteriol.*, **49**, 729–735.
- Strauss, H. (2003) Sulphur isotopes and the early Archean sulphur cycle. *Precambrian Res.*, **126**, 349–361.
- Strauss, H., Des Marais, D.J., Hayes, J.M. and Summons, R.E. (1992) Concentrations of organic carbon and maturites and elemental compositions of kerogens. In: *The Proterozoic Biosphere* (Eds J.W. Schopf and C. Klein), pp. 95–99. Cambridge University Press, Cambridge.
- Summons, R.E., Jahnke, L.L., Hope, J.M. and Logan, G.A. (1999) 2-Methylhopanoids as biomarkers for cyanobacterial oxygenic photosynthesis. *Nature*, **400**, 554–557.
- Tice, M.M. and Lowe, D.R. (2004) Photosynthetic microbial mats in the 3,416 Myr old ocean. *Nature*, **431**, 549–552.
- Trendall, A.F. (2002) The significance of iron-formation in the Precambrian stratigraphic record. *Int. Assoc. Sedimentol. Spec. Publ.*, **33**, 33–66.
- Trendall, A.F. and Blockley, J.G. (1970) The iron formations of the Precambrian Hamersley Group, Western Australia; with special reference to the associated crocidolite. *W. Austr. Geol. Surv. Bull.*, **119**, 336.
- Van Kranendonk, M.J. (2006) Volcanic degassing, hydrothermal circulation and the flourishing of early life on Earth: a review of the evidence from c. 3490–3240 Ma rocks of the Pilbara Supergroup, Pilbara Craton, Western Australia. *Earth-Sci. Rev.*, **74**, 197–240.
- Vargas, M., Kashefi, K., Blunt-Harris, E.L. and Lovely, D.R. (1998) Microbiological evidence for Fe(III) reduction on early Earth. *Nature*, **395**, 65–67.
- Waldbauer, J.R., Newman, D.K. and Summons, R.E. (2011) Microaerobic steroid biosynthesis and the molecular fossil record of Archean life. *Proc. Natl Acad. Sci. USA*, **108**, 13409–13414.
- Walker, J.C.G. (1984) Suboxic diagenesis in banded iron formations. *Nature*, **309**, 340–342.
- Widdel, F., Schnell, S., Heising, S., Ehrenreich, A., Assmus, B. and Schink, B. (1993) Ferrous iron oxidation by anoxygenic phototrophic bacteria. *Nature*, **362**, 834–836.
- Woese, C.R. (1987) Bacterial evolution. *Microbiol. Rev.*, **51**, 221–271.
- Wright, D.T. and Altermann, W. (2000) Microfacies development in Late Archaean stromatolites and oolites of

the Campbellrand Subgroup, South Africa. In: *Carbonate Platform Systems: Components and Interactions* (Eds E. Insalco, P.W. Skelton and T.J. Palmer), *Geol. Soc. London Spec. Publ.*, **178**, 51–70.

Xiong, J. (2006) Photosynthesis: what color was its origin? *Genome Biol.*, **7**(12), 245.1–245.5.

Yamaguchi, K.E., Johnson, C.M., Beard, B.L. and Ohmoto, H. (2005) Biogeochemical cycling of iron in the Archean

Palaeoproterozoic Earth: constraints from iron isotope variations in sedimentary rocks from the Kaapvaal and Pilbara Cratons. *Chem. Geol.*, **218**, 135–169.

Manuscript received 6 December 2011; revision accepted 4 April 2013