



Dr Karl Guido Rijkhoek
Director

Antje Karbe
Press Officer
Phone +49 7071 29-76788
+49 7071 29-76789
Fax +49 7071 29-5566
karl.rijkhoek[at]uni-tuebingen.de
antje.karbe[at]uni-tuebingen.de
www.uni-tuebingen.de/aktuell

Press Release

Big fish: From 5 to 150 cm in 37 million years

Oldest fossil bighead carp discovered in Vietnam

Tübingen, 12 March 2014

During excavations in the Na Duong open-cast coal mine in Vietnam, a joint team from the University of Tübingen and the Senckenberg Center for Human Evolution and Palaeoenvironment Tübingen discovered the world's oldest bighead carp. With a length of only 5 centimeters, *Planktophaga minuta* is also the smallest known fossil representative of this East Asian group. Modern bighead carp are among the largest members of the carp family, reaching a length of up to 1.5 meters and a weight of 50 kilograms.

Since 2008, an international research team led by Prof. Dr. Madelaine Böhme from the Senckenberg Center for Human Evolution and Palaeoecology (HEP) of the University of Tübingen has been studying prehistoric ecosystems and fossils in Vietnam. In the course of this research the scientists discovered approximately 37 million-year-old sediments from Lake Rhin Chua, dating to the late Eocene. These freshwater sediments contained a wealth of fossilized animals and plants.

During their studies the team discovered teeth belonging to an entirely new genus and species of fish: The oldest known bighead carp, *Planktophaga minuta*, is a representative of the East Asian group of Leuciscinae. With a length of ca. 5 centimeters it is the smallest fossil representative of this East Asian group, and a mere dwarf compared to its modern living relatives. Modern bighead carp are among the largest members of the carp family. They grow up to a length of 1.5 meters and can weigh in at 50 kilograms.

Besides *Planktophaga minuta* (which translates as "small plankton eater"), an additional six species of carp have been discovered in Lake Rhin Chua. All of them have living relatives that are still found today in China's Pearl and Yangtze River system. This is proof that the roots of the modern freshwater fish fauna in Southeast Asia reach far into the past.

Originally, the bighead carp was native to the larger rivers and stagnant water bodies of southern China. During the 1960s, bighead carp were introduced to Europe, including Germany, as a means to control aquatic plants. Only later did researchers discover that the bighead carp failed to

Please send us a copy of your article or report.

fulfill this task, since they mainly feed on animal plankton. In Europe, introduced bighead carp can be found in ponds, lakes, and occasionally in streams and rivers.

Publication Böhme, M. et al.; Na Duong (northern Vietnam) – an exceptional window into Eocene ecosystems from Southeast Asia, *Zitteliana* 53, 120–125 (2014).
Online: http://www.palmuc.de/bspg/images/pdf/10_boehme.pdf


Contact

Prof. Dr. Madeleine Böhme
Department of Geosciences
Senckenberg Center for Human Evolution and Palaeoecology
(currently away on expedition)
madelaine.boehme@senckenberg.de

Press Office

Senckenberg Gesellschaft für Naturforschung
Ilona Bröhl
Phone 069- 7542 1444
pressestelle@senckenberg.de

Press Images



Teeth of the 37 million year old bighead carp fossil
©Senckenberg

