

# Advanced Mathematical Methods

WS 2019/20

## 1 Linear Algebra

PD Dr. Thomas Dimpfl

*Chair of Statistics, Econometrics and Empirical Economics*

EBERHARD KARLS  
UNIVERSITÄT  
TÜBINGEN



WIRTSCHAFTS- UND  
SOZIALWISSENSCHAFTLICHE  
FAKULTÄT

# Outline: Linear Algebra

- 1.1 Vectors
- 1.2 Matrices
- 1.3 Special Matrices
- 1.4 Inverse of a quadratic matrix
- 1.5 The determinant
- 1.6 Calculation of the inverse
- 1.7 Linear independence and rank of a matrix

# Readings

- ▶ Knut Sydsaeter and Peter Hammond. *Essential Mathematics for Economic Analysis*. Prentice Hall, third edition, 2008 Chapters 15-16
- ▶ Knut Sydsaeter, Peter Hammond, Atle Seierstad, and Arne Strøm. *Further Mathematics for Economic Analysis*. Prentice Hall, 2008 Chapter 1

# Online Resources

MIT course on Linear Algebra (by Gilbert Strang)

- ▶ Lecture 1: Vectors, Matrices  
<https://www.youtube.com/watch?v=ZK3O402wf1c>
- ▶ Lecture 3: Multiplication and Inverse Matrices  
<https://www.youtube.com/watch?v=QVKj3LADCnA>
- ▶ Lecture 9: Independence, basis and dimension  
<https://www.youtube.com/watch?v=yjBerM5jWsc>
- ▶ Lecture 18: Properties of determinants  
<https://www.youtube.com/watch?v=srxexLishgY>

# 1.1 Vectors

## Vector operations

**multiplication** of an  $n$ -dimensional vector  $\mathbf{v}$  with a scalar  $c \in \mathbb{R}$ :

$$c \cdot \underset{(n \times 1)}{\mathbf{v}} = \begin{pmatrix} c \cdot v_1 \\ \vdots \\ c \cdot v_n \end{pmatrix}$$

**sum** of two  $n$ -dimensional vectors  $\mathbf{v}$  und  $\mathbf{w}$ :

$$\underset{(n \times 1)}{\mathbf{v}} + \underset{(n \times 1)}{\mathbf{w}} = \begin{pmatrix} v_1 + w_1 \\ \vdots \\ v_n + w_n \end{pmatrix}$$

The **difference** between two  $n$ -dimensional Vectors  $\mathbf{v}$  and  $\mathbf{w}$  is obtained by  $\mathbf{v} - \mathbf{w} = \mathbf{v} + (-1)\mathbf{w}$ .

# 1.1 Vectors

## Vector operations

Inner product (Scalar product)  $\mathbf{v}, \mathbf{w} \in \mathbb{R}^n$ :

$$\underset{(1 \times n)}{\mathbf{v}'} \cdot \underset{(n \times 1)}{\mathbf{w}} = \sum_{i=1}^n \underset{(1 \times 1)}{v_i w_i}$$

# 1.2 Matrices

## Matrix operations

**Multiplication with a scalar:**

$$\mathbf{C} = k \cdot \mathbf{A} \Leftrightarrow c_{ij} = k \cdot a_{ij} \quad \forall i, j.$$

**Addition (Subtraction) of matrices:**

for two matrices  $\mathbf{A}$  and  $\mathbf{B}$  with the same dimensions

$$\mathbf{C} = \mathbf{A} \pm \mathbf{B} \Leftrightarrow c_{ij} = a_{ij} \pm b_{ij} \quad \forall i, j.$$

# 1.2 Matrices

## Matrix multiplication

$$C = A \cdot B$$

with

$$c_{kl} = \sum_{i=1}^m a_{ki} \cdot b_{il}$$

Note: Conformity and dimensionality.

$$\begin{array}{c} \mathbf{C} \\ (n \times p) \end{array} = \underbrace{\begin{array}{c} \mathbf{A} \\ (n \times m) \end{array} \times \begin{array}{c} \mathbf{B} \\ (m \times p) \end{array}}_{\text{conformity}} \\ \underbrace{\hspace{10em}}_{\text{dimensionality}}$$



# 1.2 Matrices

## Rules of matrix multiplication

Given conformity, it holds that:

- ▶  $(A \cdot B) \cdot C = A \cdot (B \cdot C)$  (associative law)
- ▶  $(A + B) \cdot C = A \cdot C + B \cdot C$  (distributive law from the right)
- ▶  $A \cdot (B + C) = A \cdot B + A \cdot C$  (distributive law from the left)

**Power of a matrix:** For a quadratic matrix  $A$  we calculate the non-negative integer power as follows:

$$A^n = \underbrace{AA \cdots A}_{n\text{-mal}} \quad \text{with } n > 0$$

special case:  $A^0 = I$ .

## 1.2 Matrices

### Kronecker product

$\mathbf{A}$  is  $m \times n$  and  $\mathbf{B}$  is  $p \times q$ , then the Kronecker product  $\mathbf{A} \otimes \mathbf{B}$  is the  $mp \times nq$  block matrix

$$\mathbf{A} \otimes \mathbf{B} = \begin{bmatrix} a_{11}\mathbf{B} & \dots & a_{1n}\mathbf{B} \\ a_{21}\mathbf{B} & \dots & a_{2n}\mathbf{B} \\ \vdots & \ddots & \vdots \\ a_{m1}\mathbf{B} & \dots & a_{mn}\mathbf{B} \end{bmatrix}$$

## 1.2 Matrices

**Idempotent matrix:**

A quadratic matrix  $\mathbf{A}$  is idempotent if:  $\mathbf{A}^2 \equiv \mathbf{A}\mathbf{A} = \mathbf{A}$ .

**Trace of a quadratic matrix:**

$$\text{tr}(\mathbf{A}) \equiv \sum_{i=1}^n a_{ii}$$

## 1.3 Inverse of a quadratic matrix

The inverse of a matrix  $\mathbf{A}$ , expressed by  $\mathbf{A}^{-1}$ , should have the following characteristics:

$$\mathbf{A} \cdot \mathbf{A}^{-1} = \mathbf{A}^{-1} \cdot \mathbf{A} = \mathbf{I}$$

Note:

- 1.) The matrix  $\mathbf{A}$  has to be quadratic (due to conformity). Otherwise it is not invertible.
- 2.) The inverse doesn't have to exist for every single quadratic matrix
- 3.) If there is an inverse, we call the quadratic matrix *non-singular*, otherwise we call it *singular*.

## 1.3 Inverse of a quadratic matrix

4.) If there is an inverse, then it is unambiguous

Characteristics (for non-singular matrices  $A, B$ ):

- ▶  $(A^{-1})^{-1} = A$
- ▶  $(AB)^{-1} = B^{-1}A^{-1}$
- ▶  $(A')^{-1} = (A^{-1})'$

## 1.4 The determinant

### Sarrus' Rule

For a  $2 \times 2$  matrix

$$\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}$$

the determinant is defined as follows:

$$\det(\mathbf{A}) = |\mathbf{A}| = a_{11} a_{22} - a_{12} a_{21}$$

## 1.4 The determinant

An important application:

In general we can show that the determinant of a quadratic matrix with **linearly dependent columns (or rows)** has a zero determinant.

$\implies$  The determinant criterion gives us information about the linear dependency (or independency) of the rows (or rather columns) of a matrix as well as about the existence of its inverse.

## 1.4 The determinant

The determinant of the  $(3 \times 3)$ -matrix  $\mathbf{A}$  is defined as

$$\det(\mathbf{A}) = a_{11} \cdot |\mathbf{A}_{11}| - a_{12} \cdot |\mathbf{A}_{12}| + a_{13} \cdot |\mathbf{A}_{13}|$$

(cofactor formula)



## 1.4 The determinant

Illustration:

$$\mathbf{A}_{(3 \times 3)} = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}$$

Determining the **submatrices**:

Elimination of the 1<sup>st</sup> row and the 1<sup>st</sup> column of  $\mathbf{A}$  yields the **submatrix**  $\mathbf{A}_{11}$  of dimension  $(2 \times 2)$ :

$$\mathbf{A}_{(3 \times 3)} = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \implies \mathbf{A}_{11(2 \times 2)} = \begin{pmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{pmatrix}$$

## 1.4 The determinant

Elimination of the 1<sup>st</sup> row and the 2<sup>nd</sup> column of  $\mathbf{A}$  yields the submatrix  $\mathbf{A}_{12}$  of dimension  $(2 \times 2)$ :

$$\underset{(3 \times 3)}{\mathbf{A}} = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \implies \underset{(2 \times 2)}{\mathbf{A}_{12}} = \begin{pmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{pmatrix}$$

Elimination of the 1<sup>st</sup> row and the 3<sup>rd</sup> column of  $\mathbf{A}$  yields the submatrix  $\mathbf{A}_{13}$  of dimension  $(2 \times 2)$ :

$$\underset{(3 \times 3)}{\mathbf{A}} = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \implies \underset{(2 \times 2)}{\mathbf{A}_{13}} = \begin{pmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{pmatrix}$$

The determinants  $|\mathbf{A}_{ij}|$  of the submatrices  $\mathbf{A}_{ij}$  are called **subdeterminants**; They can be calculated using the *Sarrus' Rule* (if of order of 3 or lower)

## 1.4 The determinant

Alternative: Extension of the  $(3 \times 3)$ -matrix  $\mathbf{A}$  for the application of the *Rule of Sarrus*:

$$\mathbf{A}^* = \left( \begin{array}{ccc|cc} a_{11} & a_{12} & a_{13} & a_{11} & a_{12} \\ a_{21} & a_{22} & a_{23} & a_{21} & a_{22} \\ a_{31} & a_{32} & a_{33} & a_{31} & a_{32} \end{array} \right)$$

$$\begin{aligned} \det(\mathbf{A}) &= a_{11} a_{22} a_{33} + a_{12} a_{23} a_{31} + a_{13} a_{21} a_{32} \\ &\quad - a_{13} a_{22} a_{31} - a_{11} a_{23} a_{32} - a_{12} a_{21} a_{33} \end{aligned}$$

# 1.4 The determinant

## Cofactor expansion

**Calculation of the determinant for general  $n \times n$  matrices:**

Cofactor expansion *across a row  $i$* :

$$\det(\mathbf{A}) = \sum_{j=1}^n (-1)^{i+j} a_{ij} | \mathbf{A}_{ij} |$$

Alternatively: Cofactor expansion *down a column  $j$* :

$$\det(\mathbf{A}) = \sum_{i=1}^n (-1)^{i+j} a_{ij} | \mathbf{A}_{ij} |$$

Note: The product  $(-1)^{i+j} | \mathbf{A}_{ij} |$  is called **cofactor**.

# 1.4 The determinant

## Properties of determinants

for  $\mathbf{A}$  and  $\mathbf{B}$  with dimension  $n \times n$ :

- 1.) The exchange of two rows or two columns of a matrix leads to a change in the sign of the determinant.
- 2.) The determinant doesn't change its value if we add to a row (column) within a matrix the multiple of another row (column).
- 3.) The determinants of a matrix and its transpose are equal:

$$\det(\mathbf{A}) = \det(\mathbf{A}')$$

- 4.) Multiplying all components of a ( $n \times n$ ) matrix with the same factor  $k$  leads to a change in the value of the determinant by the factor  $k^n$ :

$$\det(k\mathbf{A}) = k^n \det(\mathbf{A})$$

# 1.4 The determinant

## Properties of determinants

- 5.) The determinant of every identity matrix is equal to 1; the determinant of every zero matrix is equal to 0.
- 6.) The determinant of the product of  $\mathbf{A}$  and  $\mathbf{B}$  equals the product of the determinants of  $\mathbf{A}$  and  $\mathbf{B}$ :

$$\det(\mathbf{A} \cdot \mathbf{B}) = \det(\mathbf{A}) \cdot \det(\mathbf{B})$$

- 7.) From 6.) follows for a regular matrix  $\mathbf{A}$  that:

$$\det(\mathbf{A}^{-1}) = \frac{1}{\det(\mathbf{A})}$$

- 8.) In general:  $\det(\mathbf{A} + \mathbf{B}) \neq \det(\mathbf{A}) + \det(\mathbf{B})$ .

## 1.5 Calculation of the inverse

We can determine regularity/non-singularity/invertibility of the *square matrix*  $\mathbf{A}$  using the determinant. It holds that

$$\det(\mathbf{A}) \neq 0 \Leftrightarrow \mathbf{A}^{-1} \text{ exists.}$$

## 1.5 Calculation of the inverse

In general: The inverse of the  $(n \times n)$ -matrix  $\mathbf{A}$  is denoted as

$$\mathbf{A}^{-1} = \mathbf{B} = \begin{pmatrix} b_{11} & \dots & b_{1n} \\ \vdots & & \vdots \\ b_{n1} & \dots & b_{nn} \end{pmatrix}$$

We get every single element of  $\mathbf{B}$  by

$$b_{ij} = \frac{1}{|\mathbf{A}|} (-1)^{(i+j)} |\mathbf{A}_{ji}| \quad (\text{note the index!})$$

In order to get the element  $b_{ij}$ , you have to calculate the subdeterminant  $\mathbf{A}_{ji}$  crossing out the  $j$ -th row and the  $i$ -th column of  $\mathbf{A}$ .



## 1.6 Linear independence and rank of a matrix

### Linear combination of vectors

#### Definition: linear combination

For the vectors  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k \in \mathbb{R}^n$  a  $n$ -dimensional vector  $\mathbf{w}$  is called **linear combination** of vectors  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ , if there are real numbers  $c_1, c_2, \dots, c_k \in \mathbb{R}$ , such that:

$$\mathbf{w} = c_1 \cdot \mathbf{v}_1 + c_2 \cdot \mathbf{v}_2 + \dots + c_k \cdot \mathbf{v}_k = \sum_{i=1}^k c_i \cdot \mathbf{v}_i .$$

# 1.6 Linear independence and rank of a matrix

## Linear independence

### Definition: linear independence

The vectors  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k \in \mathbb{R}^n$  are called **linearly independent**, if

$$c_1 \cdot \mathbf{v}_1 + c_2 \cdot \mathbf{v}_2 + \dots + c_k \cdot \mathbf{v}_k = \mathbf{0} \quad \text{with} \quad c_1, c_2, \dots, c_k \in \mathbb{R}$$

is only attainable with  $c_1 = c_2 = \dots = c_k = 0$ . Otherwise they are called **linearly dependent** and  $\mathbf{v}_1 = d_2 \cdot \mathbf{v}_2 + \dots + d_k \cdot \mathbf{v}_k$  (with  $d_2, d_3, \dots, d_k \in \mathbb{R}$ ) applies.

## 1.6 Linear independence and rank of a matrix

### Rank

The **rank** of the  $n \times m$ -matrix  $\mathbf{A}$  ( $\text{rk}(\mathbf{A})$ ) is determined by the maximum number of linearly independent columns (rows) of the matrix  $\mathbf{A}$ .

$$\text{rk}(\mathbf{A}) \leq \min(m, n)$$

For every matrix the column rank equals the row rank.

The rank criterion allows to determine whether a quadratic  $n \times n$  matrix  $\mathbf{A}$  is regular/non-singular or not:

$$\text{rk}(\mathbf{A}) = n \Rightarrow \textit{non - singular}$$

$$\text{rk}(\mathbf{A}) < n \Rightarrow \textit{singular}$$

## 1.6 Linear independence and rank of a matrix

### Properties of the rank

- 1.) The rank of a matrix doesn't change if you exchange rows or columns among themselves.
- 2.) The rank of a matrix  $\mathbf{A}$  is equal to the rank of the transpose  $\mathbf{A}'$ .
- 3.) For a  $(m \times n)$  matrix  $\mathbf{A}$  the following applies:  
 $\text{rk}(\mathbf{A}) = \text{rk}(\mathbf{A}'\mathbf{A})$ , whereby  $\mathbf{A}'\mathbf{A}$  is quadratic.

## 1.6 Linear independence and rank of a matrix

### Determination of the rank of a matrix

- 1.) We consider all quadratic submatrices of a matrix of which the determinants are not 0. Then we search for the determinant of highest order. The order of this determinant is equal to the rank of the matrix.
- 2.) Using gaussian algorithm
- 3.) Using eigenvalues