



Press Release

When hornless rhinos lived in Europe

International team analyzes fossil skulls and redefines species: relatives of today's rhinos had no horn and died out five million years ago

Tübingen, 02.11.2023

Paleontologists from Tübingen have redefined a rhinoceros genus that had fallen into oblivion: *Eochilotherium* lived more than five million years ago and did not have a horn on its nose. Hornless rhinos were known to be ancestors of today's species. An international research team from Germany, Greece, Bulgaria and South Africa shows in the *Journal of Vertebrate Paleontology* that these were more diverse than previously thought. Panagiotis Kampouridis of the Senckenberg Centre for Human Evolution and Palaeoenvironment at the University of Tübingen re-examined the fossil skulls of hornless rhinos.

Today's rhinos bear the characteristic horns on their nose and/or forehead and live Africa and Asia. Three of the five species are currently threatened with extinction. In the 40-million-year-old evolutionary history of large herbivores, there have been numerous species that have gone extinct, including many without horns.

Hornless rhinos were perhaps the most diverse group of the entire family and one of the most species-rich genera is known as *Chilotherium*. They lived in Asia as well as in Eastern and Southeastern Europe. They were smaller than today's species and with extremely short legs were probably not as eager to run as today's rhinos. They had tusk-like incisors and grazed in open landscapes. Five million years ago at the latest chilothers became extinct in Europe and a little later also in Asia, probably also because of changed climatic conditions.

The originals of two chilothere species, *Chilotherium schlosseri* and *Eochilotherium samium* – so called holotypes –, were destroyed during the Second World War. This made the identification of the two species difficult. Kampouridis, a PhD student and first author of the publication, therefore searched for new evidence, traveling to various European museums.

University of Tübingen
Public Relations Department

Dr. Karl Guido Rijkhoek
Director

Antje Karbe
Press Officer

Phone +49 7071 29-76788
+49 7071 29-77853
karl.rijkhoeck[at]uni-tuebingen.de
antje.karbe[at]uni-tuebingen.de

Senckenberg Nature Research
Society
Press Office

Judith Jördens
Phone +49 69 7542 1434
judith.joerdens[at]senckenberg.de

pressestelle[at]senckenberg.de
www.senckenberg.de/presse

Using two skulls, from the Museum der Natur Hamburg and from the Senckenberg Naturmuseum Frankfurt, he finally succeeded in redefining the species. The skulls confirm that there was another genus of hornless rhinoceroses besides *Chilotherium*, *Eochilotherium*. This is shown by the shape of the skull and teeth - *Eochilotherium samium* has a slightly smaller and narrower skull and less pronounced enamel folds in the upper teeth than *Chilotherium schlosseri*.

A complex history of migration

The authors of the study even assume that two other species from China, which, like *Eochilotherium samium*, were previously included in the genus *Chilotherium*, are actually more closely related to *Eochilotherium* or may even represent another genus. "This changes our understanding about the chilothere group, which was previously thought to contain only two genera. According to the present results, there are three, possibly four," says co-author Professor Nikolai Spassov of the Bulgarian Academy of Sciences.

"Our results also provide insight into the biogeography of the short-legged chilotheres, which, according to previous findings, migrated to Europe from Asia at least twice. However, it is possible that their distribution history is even more complex," says Professor Madelaine Böhme of the University of Tübingen and the Senckenberg Center for Human Evolution and Palaeoenvironment.

This fundamental study is therefore the basis of a detailed study of the last European chilotheres, which did not survive despite high diversity, say the authors. It shows the importance of detailed study of species that are already known, they add. "Rhinos, like other large herbivores, have very important roles in their respective ecosystems, and their loss can lead to far-reaching consequences for the rest of the fauna," Kampouridis says. "This is as true for fossil ecosystems as it is for present-day ecosystems."


Publication:

Panagiotis Kampouridis, Georgia Svorligkou, Nikolaos Kargopoulos, Nikolai Spassov & Madelaine Böhme: Revision of the late Miocene hornless rhinocerotods from Sanmos Island (Greece) with the designation of neotypes and implications for the European chilotheres. *Journal of Vertebrate Paleontology*, <https://doi.org/10.1080/02724634.2023.2254360>


Contact:

Panagiotis Kampouridis
University of Tübingen
Geosciences – Terrestrial Palaeoclimatology
Phone +49 7071 29-73561
panagiotis.kampouridis@uni-tuebingen.de

Professor Dr. Madelaine Böhme
University of Tübingen
Geosciences – Terrestrial Palaeoclimatology and
Senckenberg Center for Human Evolution and Palaeoenvironment
Phone +49 7071 29-73191
m.boehme@ifg.uni-tuebingen.de



On the basis of these skulls, the relationships of the hornless rhinoceros species were redefined. Photos: P. Kampouridis



3D surface models of the examined skulls of hornless rhinoceros species. Photos: P. Kampouridis