

EBERHARD KARLS  
UNIVERSITÄT  
TÜBINGEN



WIRTSCHAFTS- UND  
SOZIALWISSENSCHAFTLICHE  
FAKULTÄT

Chair of Statistics, Econometrics and Empirical Economics

PD Dr. Thomas Dimpfl

**S414**  
**Advanced Mathematical Methods**  
Exercises

WS 2019/20

## LINEAR ALGEBRA

EXERCISE 1 **Quadratic Form**

Given the matrix

$$\mathbf{A} = \begin{pmatrix} 2 & -2 \\ 1 & 2 \end{pmatrix} .$$

- Determine the definiteness of the quadratic form  $Q = \mathbf{x}'\mathbf{A}\mathbf{x}$ .
- Explain in two sentences maximum what this means for the graph  $\{(x_1, x_2, Q) | Q = (x_1; x_2)\mathbf{A}(x_1; x_2)'\}$ .

EXERCISE 2 **Quadratic Form**

Write the quadratic form

$$Q = 4x_1^2 + 4x_1x_2 - x_2^2$$

in matrix notation and determine its definiteness.

**EXERCISE 3 Sign definiteness**

Express each quadratic form below as a matrix product involving a *symmetric* coefficient matrix:

a)  $q = 3u^2 - 4uv + 7v^2$

b)  $q = u^2 + 7uv + 3v^2$

c)  $q = 8uv - u^2 - 31v^2$

d)  $q = 6xy - 5y^2 - 2x^2$

e)  $q = 3u_1^2 - 2u_1u_2 + 4u_1u_3 + 5u_2^2 + 4u_3^2 - 2u_2u_3$

f)  $q = -u^2 + 4uv - 6uv - 4v^2 - 7w^2$

**EXERCISE 4 Sign definiteness**

Given a quadratic form  $u'Du$ , where  $D$  is  $2 \times 2$ , the characteristic equation of  $D$  can be written as:

$$\begin{vmatrix} d_{11} - r & d_{12} \\ d_{21} & d_{22} - r \end{vmatrix} = 0 \quad (d_{12} = d_{21})$$

Expand the determinant; express the roots of this equation by use of the quadratic formula and deduce the following:

a) No imaginary number (a number involving  $\sqrt{-1}$ ) can occur in  $r_1$  and  $r_2$ .

b) To have repeated roots, the matrix  $D$  must be in the form of  $\begin{pmatrix} c & 0 \\ 0 & c \end{pmatrix}$

c) To have either positive or negative definiteness, the determinant of the matrix  $D$  must vanish, i.e.  $|D| = 0$ .

**Solution Exercise 1:**

a) positive definite

**Solution Exercise 2:**

$$\mathbf{Q} = \mathbf{x}'\mathbf{A}\mathbf{x} \text{ with } A = \begin{pmatrix} 4 & 2 \\ 2 & -1 \end{pmatrix}$$

 $\mathbf{A}$  is indefinite**Solution Exercise 3:**Quadratic form:  $q = \mathbf{x}'\mathbf{A}\mathbf{x}$ 

a)

$$q = \begin{pmatrix} -u \\ v \end{pmatrix}' \begin{pmatrix} 3 & 2 \\ 2 & 7 \end{pmatrix} \begin{pmatrix} -u \\ v \end{pmatrix}$$

b)

$$q = \begin{pmatrix} u \\ v \end{pmatrix}' \begin{pmatrix} 1 & 3.5 \\ 3.5 & 3 \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix}$$

c)

$$q = \begin{pmatrix} u \\ v \end{pmatrix}' \begin{pmatrix} -1 & 4 \\ 4 & -31 \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix}$$

d)

$$q = \begin{pmatrix} x \\ y \end{pmatrix}' \begin{pmatrix} -2 & 3 \\ 3 & -5 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$$

e)

$$q = \begin{pmatrix} u_1 \\ -u_2 \\ u_3 \end{pmatrix}' \begin{pmatrix} 3 & 1 & 2 \\ 1 & 5 & 1 \\ 2 & 1 & 4 \end{pmatrix} \begin{pmatrix} u_1 \\ -u_2 \\ u_3 \end{pmatrix}$$

f)

$$q = \begin{pmatrix} u \\ v \\ -w \end{pmatrix}' \begin{pmatrix} -1 & 2 & 3 \\ 2 & -4 & 0 \\ 3 & 0 & -7 \end{pmatrix} \begin{pmatrix} u \\ v \\ -w \end{pmatrix}$$