

Size-Dependent Reduction Kinetics of Iron Oxides in Single and Mixed Mineral Systems

Xiyang Xu, Muammar Mansor, Guoxiang Li, Tsz Ho Chiu, Stefan B. Haderlein, Andreas Kappler, and Prachi Joshi*



Cite This: <https://doi.org/10.1021/acs.est.4c08032>



Read Online

ACCESS |


Metrics & More

Article Recommendations

Supporting Information

ABSTRACT: Iron(III) (oxyhydr)oxide minerals with varying particle sizes commonly coexist in natural environments and are susceptible to both chemical and microbial reduction, affecting the fate and mobility of trace elements, nutrients, and pollutants. The size-dependent reduction behavior of iron (oxyhydr)oxides in single and mixed mineral systems remains poorly understood. In this study, we used microbial and mediated electrochemical reduction approaches to investigate the reduction kinetics and extents of goethite and hematite. We found that small particles were preferentially reduced relative to their large counterparts in single and mixed mineral systems regardless of microbial or electrochemical treatments, which is attributed to the combined effect of higher thermodynamic favorability and greater surface availability. In mixed mineral systems, small particles were reduced slightly faster, whereas large particles were reduced notably slower and less extensively than solely predicted from single mineral systems. Specifically, when reduced alone, small particles showed Fe(III) reduction rate constants that were 1.5- to 3.6-fold higher than large particles, while when reduced together, the reduction rate constants for small particles were 6- to 21-fold higher than the rate constants for large particles. These collective findings provide new insights into the pivotal role of nanoparticulate iron (oxyhydr)oxides in environmental redox reactions.

KEYWORDS: iron minerals, nanoparticles, microbial reduction, mediated electrochemical reduction, reduction kinetics



INTRODUCTION

Iron(III) (oxyhydr)oxide minerals, such as goethite (α -FeOOH) and hematite (α -Fe₂O₃), are ubiquitous in soils and sediments,^{1–3} where they span a continuum of sizes, from nanometer to the millimeter scale.^{2,4} Nanoparticles behave remarkably differently from their larger counterparts, resulting in enhanced roles in the biogeochemical cycling and fate of nutrients, trace elements, and contaminants via redox reactions.^{5–11} For instance, nanoparticulate iron (oxyhydr)oxides can act as terminal electron acceptors in anaerobic microbial respiration,^{12–15} resulting in their dissolution and the subsequent release of associated trace elements.^{16,17}

The significance of iron redox reactions has led to extensive research over the past decades into the susceptibility of iron (oxyhydr)oxides toward microbially and chemically driven reduction processes. The extents and rates of Fe(III) mineral reduction have been suggested to depend on the characteristics of the minerals such as particle size, morphology, crystallinity, reactive surface area, vacancies, solubility, dissolution, thermodynamic properties as well as cell–mineral attachment, secondary biomineralization, and presence of electron shuttles.^{14,18–30} Notably, Roden (2003) observed a strong linear correlation between the initial rate of microbial Fe(III)

reduction and mineral surface area and suggested that microbial Fe(III) mineral reduction was not significantly influenced by the thermodynamic properties of the iron (oxyhydr)oxides.²⁰ In contrast, Bonneville et al. (2004, 2006) proposed that solubility and coverage of the cell surface by hematite nanoparticles were rate-controlling factors in both biotic and abiotic reduction of the hematite.^{25,26} Yan et al. (2008) suggested that the reduction rates of nanohematite were likely to be proportional to the cell–mineral contact area and not to the total surface area.²⁷ In addition, Aeppli et al. (2018, 2022) postulated that thermodynamics controlled goethite and hematite reduction kinetics in mediated electrochemical experiments.^{28,29} In these studies as well as other past work, identifying effects of particle size on reduction rates and extents has been elusive because comparative experiments typically investigated either different minerals or the same

Received: August 2, 2024

Revised: January 15, 2025

Accepted: January 16, 2025

mineral with one or two properties that were varied. However, evaluating these variables independently is difficult, as they are all related. Therefore, accurately determining the size effects on reduction kinetics of nanoparticles requires systematic investigation of the same mineral with all relevant explanatory factors.

While abundant literature has described the reduction of individual iron (oxyhydr)oxides, these minerals often coexist as mixtures of mineral phases^{1,31–35} and varying particle sizes in the environment.^{36,37} For instance, coexisting goethite and hematite have been reported in ocean sediments,³² lacustrine sediments,³³ loess-paleosol sequences,³⁴ and soils.^{31,35} Although the coexistence of minerals in the environment is expected, there have been only a few studies examining how coexistence affects overall reactivity.^{38–41} For example, Notini et al. (2019) observed that goethite with defects displayed an initial microbial Fe(III) reduction rate about 6-fold higher than goethite with fewer defects, when these minerals were present together.³⁸ Hua et al. (2022) studied microbial Fe(III) reduction with respect to the coexistence of hematite with multiple facets, showing that hematite nanoplates were reduced 1.8 times more and 1.6 times faster than hematite nanorods.³⁹ Other studies have looked at processes other than reduction in mixed systems: Notini et al. (2023) demonstrated that the coexistence of goethite promoted the transformation of ferrihydrite into goethite.⁴⁰ Higher ratios of goethite to ferrihydrite have been shown to promote heteroaggregation, which may impact their combined reactivity.⁴¹ However, no studies to date have examined the reduction behavior of iron (oxyhydr)oxides in mixed mineral systems with varying particle sizes.

In the present study, the reduction kinetics and extent of iron (oxyhydr)oxides in single and mixed mineral systems were systematically investigated using microbial and mediated electrochemical reduction (MER) approaches. The work had two specific objectives. (i) To evaluate the impact of mineral-specific related variables on their reduction kinetics and extents within single mineral systems through redundancy analysis (RDA) and Pearson correlation analysis. Here, we utilized the unique capability of MER to adjust the thermodynamic boundary conditions in order to study the thermodynamic driving force of iron (oxyhydr)oxide reduction, complementary to microbial reduction.^{42,43} (ii) To determine the size preferential reduction behavior of iron (oxyhydr)oxides within environmentally relevant mixed mineral systems. We employed Fe isotope-labeled minerals to track the origin of reduced Fe in mixed mineral systems during microbial reduction processes. The findings of this research enhance our understanding of the reduction susceptibility of iron (oxyhydr)oxides with varying particle sizes in the environments.

MATERIALS AND METHODS

Iron (oxyhydr)oxide Synthesis and Characterization.

Goethite and hematite with varied particle sizes were synthesized based on previous methods.^{44–46} The synthesized minerals are termed Gt_2000, Gt_90, Gt_30, Hm_300, Hm_40, and Hm_8 based on their particle size. We synthesized a second set of iron (oxyhydr)oxides for mixed mineral experiments, which included naturally abundant Fe minerals (referred to as ^{NA}mineral, containing 5.82% of ⁵⁴Fe) and ⁵⁶Fe-enriched Fe minerals (referred to as ⁵⁶mineral, containing negligible ⁵⁴Fe, prepared from ⁵⁶Fe-enriched Fe⁰ (Isoflex, 99.94% purity)). The purities of the synthesized iron

(oxyhydr)oxides were determined using micro-X-ray diffraction (μ -XRD) and ⁵⁷Fe Mössbauer spectroscopy. The morphologies and particle size distributions of the iron (oxyhydr)oxides were characterized using scanning electron microscopy (SEM) and transmission electron microscopy (TEM). The BET surface area was determined by N₂ sorption at 77 K. The zeta potentials and aggregation sizes of the iron (oxyhydr)oxides were measured by dynamic light scattering (DLS, Malvern Zetasizer Nano ZS).⁴⁷ Details regarding the synthesis and characterization are provided in the SI, Section S1.

Mediated Electrochemical Fe(III) Reduction. To study reduction kinetics and extents of the iron (oxyhydr)oxides at different applied reduction potentials ($E_{\text{H}}^{\text{MER}}$) as a function of particle size, mediated electrochemical reduction (MER) experiments were conducted under anoxic conditions inside a glovebox (100% N₂ atmosphere, O₂ < 2 ppm). The electrochemical setup consisted of 9 mL glassy carbon cells as the working electrode (WE), Ag/AgCl as the reference electrode, and a platinum wire as the counter electrode. The cells were controlled by eight-channel potentiostats (model 1000B, CH Instruments). The $E_{\text{H}}^{\text{MER}}$ was measured against Ag/AgCl reference electrodes, but we herein report values versus standard hydrogen electrode (SHE). Throughout the experiments, the solution in each WE cell was stirred continuously with a glass-coated stir bar.

To perform the MER, we first filled the WE cells to 6.5 mL with buffer solution (HEPES, 10 mM, pH 7) with NaCl as an electrolyte (100 mM) and then applied a defined and constant $E_{\text{H}}^{\text{MER}}$ to the WE. After the background current had stabilized at a low value (<4 μ A), mediator (10 mM) was added to the cell: 9,10-anthraquinone-2,6-disulfonic acid disodium salt (AQDS, $E_{\text{H}}^0 = -0.18$ V)⁴⁸ at $E_{\text{H}}^{\text{MER}} = -0.25$ V, diquat (DQ, $E_{\text{H}}^0 = -0.35$ V)⁴⁹ at $E_{\text{H}}^{\text{MER}} = -0.35$ V, and 4,4'-bipyridinium-1,1'-bis(2-ethylsulfonate) (ZiV, $E_{\text{H}}^0 = -0.41$ V)⁴³ at $E_{\text{H}}^{\text{MER}} = -0.53$ V. Each mediator was transferred into the WE cell in duplicate additions, followed by triplicate spikes of the iron (oxyhydr)oxides and finally duplicate additions of the mediator (SI, Figure S6). The intervals between spikes were long enough to allow the current to stabilize (<4 μ A). The experiments were designed such that the concentration of reduced mediator (833 μ M) was approximately 6 times higher than the iron(III) concentration (138 μ M) after the first addition of the iron (oxyhydr)oxide suspension. At this ratio, maximum iron (oxyhydr)oxide reduction rates have been reported to be independent of the mediator concentration.²⁸

The reductive current peaks in response to the four mediator additions were utilized to determine the rate constants of the mediator reduction in individual cells. These rate constants were subsequently employed to normalize the iron (oxyhydr)oxide reduction rate constants calculated from individual cells (see the Data Analysis section below).

Microbial Fe(III) Reduction. To study the kinetics and extents of microbial reduction of iron (oxyhydr)oxides, we performed cell suspension experiments using the fermentative Fe(III)-reducer *Shewanella oneidensis* MR-1 (details of incubation in SI, Section S3). Batch microbial Fe(III) reduction experiments were carried out in triplicates using serum bottles filled with 25 mL of anoxic buffer solution (HEPES 15 mM, pH 7). The HEPES buffer was chosen to avoid interactions with dissolved Fe(II) and minimal effect on nanoparticle dissolution or etching.⁵⁰ Sodium lactate (5 mM) and iron (oxyhydr)oxides (10 mM) were used as electron

donors and acceptors, respectively, and AQDS (20 μM) was employed as an electron shuttle to mimic natural redox-active mediators such as NOM and EPS. Washed cells with a concentration of 5×10^8 cells/mL were added into the reactors. The cell number was determined using optical density (OD) measurements. Serum bottles were placed (stationary) at 28 °C in the dark for 7 days. An aliquot of the suspension (0.5 mL) was periodically removed (at 3, 6, 12, 25, 35, 50, and 75 h) inside the N_2 -filled glovebox, with the aqueous and solid phases separated by centrifugation (12,100g, 15 min). The solids were sequentially extracted using 0.5 M HCl (30 min at 25 °C in the glovebox) and 6 M HCl (8 h at 60 °C in the glovebox). Based on the above sampling, we quantified the concentration of aqueous, adsorbed, and total Fe(II) using the ferrozine assay.^{51,52}

Microbial Fe(III) Reduction of Isotopically Labeled Mixed Minerals. To study size preferential microbial reduction in environmentally relevant complex systems with varying particle sizes and mineral types, we designed three experimental treatments: a mix of (1) $^{56}\text{Gt}_{2000}$ (Fe concentration of 5 mM) and $^{\text{NA}}\text{Gt}_{90}$ (5 mM), (2) $^{56}\text{Hm}_{40}$ (3.5 mM) and $^{\text{NA}}\text{Hm}_8$ (6.5 mM), and (3) $^{\text{NA}}\text{Gt}_{90}$ (4.5 mM) and $^{56}\text{Hm}_8$ (5.5 mM). The concentrations were chosen so as to approximate their number-based size distribution based on Pareto's law, which shows that particle number concentrations increase logarithmically with each order of magnitude decrease in particle size.^{36,37} Detailed calculations are provided in the SI, Section S5. The use of mixtures of $^{\text{NA}}$ mineral and 56 mineral allowed us to utilize ^{54}Fe and ^{56}Fe to track the origin of the reduced iron.

Experiments were performed as described above for microbial Fe(III) reduction. The isotopic compositions of obtained aqueous phase and sequentially HCl-extracted phases were determined by inductively coupled plasma mass spectrometry (ICP-MS, Agilent 7900), with standard deviations <10% for both ^{54}Fe and ^{56}Fe .

Data Analysis. The number of electrons accepted by the iron (oxyhydr)oxides in MER experiments, q [mol_e^-], was quantified by integration of the reductive current in response to iron (oxyhydr)oxide spikes, according to eq 1.⁴⁹

$$q = \frac{1}{F} \int_{t_0}^{t_{\text{end}}} I_{\text{red}} dt \quad (1)$$

where I_{red} [A] is the background-subtracted reductive current, F is the Faraday constant, and t_0 and t_{end} [s] represent the initial and end integration boundaries for each current peak. Background subtraction and peak area integration were manually performed using Origin 2021 (9.8).

- (a) The observed rate constants for MER reduction, k_{obs} [s^{-1}], were determined from the current responses to the first addition of iron (oxyhydr)oxides in MER experiments, according to eq 2.⁴³

$$-k_{\text{obs}} \cdot t = \ln \left(\frac{\text{Fe}_t^{3+}}{\text{Fe}_{t_0}^{3+}} \right) = \ln \left(1 - \frac{\frac{1}{F} \int_{t_0}^t I(t) dt}{q_{\text{total}}} \right) \quad (2)$$

where t_0 and t [s] are the time of iron (oxyhydr)oxide addition and the time during (oxyhydr)oxide reduction, respectively; $\text{Fe}_t^{3+}/\text{Fe}_{t_0}^{3+}$ denotes the fraction of the remaining ferric iron at time t relative to initially added ferric iron in the WE cell; and $I(t)$ [A] is the

background-subtracted reductive current. k_{obs} was determined from the slope of a linear regression line fitted to the reductive current response plotted as $\ln(\text{Fe}_t^{3+}/\text{Fe}_{t_0}^{3+})$ versus time, where data were collected from the time point at maximum reductive current to the time point at 90% of final reduction extent. We considered only the first addition due to its greater reduction extent and lower activities of Fe^{2+} relative to the second and third additions.

The reduction rate constants for the mediator, $k_{\text{obs}}^{\text{med}}$ [s^{-1}], were determined from the reductive peak in response to mediator additions using a modified eq 2 (i.e., $\ln(\text{med}_{\text{red}}(t)/\text{med}_{\text{red}}(t_0))$ instead of $\ln(\text{Fe}_t^{3+}/\text{Fe}_{t_0}^{3+})$). $k_{\text{obs}}^{\text{med}}$ was utilized to normalize differences between separate WE cells, which might result from the stirring speed and depth of the counter electrode in the cell (eq 3).

$$k_{\text{obs}}^* = k_{\text{obs}} \cdot \frac{k_{\text{obs,av}}^{\text{med}}}{k_{\text{obs}}^{\text{med}}} \quad (3)$$

where k_{obs}^* [s^{-1}] is the corrected observed rate constant for iron (oxyhydr)oxide reduction and $k_{\text{obs,av}}^{\text{med}}$ [s^{-1}] is the averaged mediator reduction rate constants for each mediator across all experiments.

- (b) The microbial reduction rate was calculated based on pseudo-first-order kinetics. The rate constant, k' , was determined from the slope of a linear regression plotted as $\ln(\text{Fe}_t^{3+}/\text{Fe}_{\text{total}}^{3+})$ versus time, according to eq 4.¹⁸

$$-k' \cdot t = \ln \left(\frac{\text{Fe}_t^{3+}}{\text{Fe}_{\text{total}}^{3+}} \right) = \ln \left(1 - \frac{\text{Fe}_t^{2+}}{\text{Fe}_{\text{total}}^{3+}} \right) \quad (4)$$

where t [h] is the time, $\text{Fe}_t^{3+}/\text{Fe}_{\text{total}}^{3+}$ denotes fraction of the remaining ferric iron at time t relative to initially added ferric iron, and Fe_t^{2+} is the ferrous iron concentration at time t , which was determined following 6 M HCl extraction.

Thermodynamic Calculations.

- (a) Standard reduction potential (E_{H}^0) of iron (oxyhydr)oxides was calculated from standard Gibbs free energy of reaction (ΔG_{rxn}^0). The Gibbs free energy of nanoparticle formation $\Delta G_{\text{f,i}}^0$ was calculated based on the published values for bulk minerals and corrected for increased surface energy of nanoparticles through published mineral-specific enthalpies of hydrated surfaces (ΔH_s^h) and molar volumes (S_m), assuming geometric surface area (SA) and volume (V), according to eqs 5 and 6.^{53–55}

$$E_{\text{H}}^0 = -\frac{\Delta G_{\text{rxn}}^0}{nF} = -\frac{\left(\sum v_i \Delta G_{\text{f,i}}^0 \right)_{\text{products}} - \left(\sum v_i \Delta G_{\text{f,i}}^0 \right)_{\text{reactants}}}{nF} \quad (5)$$

$$\Delta G_{\text{f,i}}^0 = \Delta G_{\text{f,bulk}}^0 + \text{SA} \cdot \frac{S_m}{V} \cdot \Delta H_s^h \quad (6)$$

where $\Delta G_{\text{f,bulk}}^0$ [kJ/mol] is the Gibbs free energy of bulk mineral formation of reactants and products, n denotes the number of transferred electrons, and v_i represents the stoichiometric coefficient of species i in the balanced chemical equations (SI, Section S4).

- (b) Reaction free energies, $\Delta_r G$ [kJ/mol], in MER experiments were calculated from differences in reduction potential between iron (oxyhydr)oxide and applied potential, according to eq 7.

$$\Delta_r G = -nF \cdot (E_H^{\text{oxide}} - E_H^{\text{MER}}) \quad (7)$$

E_H^{oxide} was calculated from the Nernst equation according to eq 8.

$$E_H^{\text{oxide}} = E_H^0 - \frac{RT}{nF} \ln\{\text{Fe}_{\text{aq}}^{2+}\} + 3 \frac{RT}{nF} \ln\{\text{H}^+\} \quad (8)$$

where R is the ideal gas constant, T is the absolute temperature, and $\{\text{Fe}_{\text{aq}}^{2+}\}$ denotes the activity of aqueous ferrous iron calculated from Davies equation.⁵⁶ According to eqs 7 and 8, $\Delta_r G$ is dependent on the dissolved ferrous iron activity. We herein calculated $\Delta_r G$ for selected $\{\text{Fe}_{\text{aq}}^{2+}\}$ that corresponded to the time of the maximum rate of electron transfer, when data collection began to calculate rate constant k_{obs} . A detailed description of the $\Delta_r G$ calculations is provided in the SI, Section S4.

Statistical Analysis. Detrended correspondence analysis (DCA) and redundancy analysis (RDA) were conducted using CANOCO 5 software (Microcomputer Power Co., USA), and Pearson correlation was performed with Origin 2021 (9.8) (OriginLab Co., USA). RDA, widely used in ecology for direct gradient analysis, can evaluate the relationship between a set of explanatory and the target variables, irrespective of whether the variables are related.⁵⁷ Before RDA, detrended correspondence analysis (DCA) was performed to determine whether the data sets were appropriate for RDA. If the gradient lengths of the four axes calculated from DCA are less than 3, then RDA was applicable. The DCA results for the variables in both the MER and microbial reduction experiments indicate gradient lengths of 0.18 and 0.28, respectively, suggesting that the data set was suitable for RDA.

RESULTS AND DISCUSSION

Characterization of Goethite and Hematite Nanoparticles. To study the size-dependent reduction kinetics and extents of goethite and hematite in single and mixed mineral systems, we synthesized goethite and hematite with a range of different sizes (30–2000 nm for goethite and 8–300 nm for hematite). The μ -XRD patterns (SI, Figure S1) and Mössbauer spectra (SI, Figure S2 and Table S1) indicated pure goethite and hematite. Widths of the reflections in XRD patterns increased with decreasing sizes, indicating a lower crystallinity. Additionally, differences in Mössbauer fitting parameters showed small variations in magnetic properties of samples. Based on the average length of goethite particles and mean diameters of hematite particles in TEM or SEM images (SI, Figures S3–S5), the minerals are referred to heretofore as Gt_2000, Gt_90, Gt_30, Hm_300, Hm_40, and Hm_8.

Mediated Electrochemical Fe(III) Reduction of Goethite and Hematite. To investigate size-dependent reduction of goethite and hematite via a mediated electrochemical approach, we measured reductive current responses to additions of these two iron (oxyhydr)oxides in electrochemical cells at different applied potentials that contained an electron transfer mediator ($E_H^{\text{MER}} = -0.53, -0.35, -0.25$ V vs SHE at pH 7.0). Figure 1 shows reductive current responses to additions of goethite (Figure 1 A) and hematite (Figure 1 B) at




Figure 1. Exemplary reductive current responses to the addition of goethite (Gt) (A) and hematite (Hm) (B) to electrochemical cells at pH 7.0 and $E_H^{\text{MER}} = -0.35$ V. Gt_2000, Gt_90, and Gt_30 denote goethite sizes of 2000, 90, and 30 nm, respectively. Hm_300, Hm_40, and Hm_8 denote hematite sizes of 300, 40, and 8 nm, respectively. Colored areas represent the number of electrons transferred by peak integration, and the percentage values represent the total Fe(III) reduction extent.

$E_H^{\text{MER}} = -0.35$ V. We observed that decreasing particle sizes resulted in increasing heights and areas of reductive current peaks for goethite and hematite. Similar trends were observed at other E_H^{MER} values (-0.53 and -0.25 V) (SI, Figures S2, S7–S9).

We determined the reduction extents and rates from the reductive current responses. At $E_H^{\text{MER}} = -0.53$ V, the reduction extents exceeded 90% for both goethite and hematite regardless of particle size (SI, Table S2), consistent with the thermodynamic favorability of complete iron (oxyhydr)oxide reduction at this low E_H .²⁸ By contrast, the reduction extents were lower with increasing (i.e., less negative) E_H^{MER} of -0.35 and -0.25 V. Smaller-sized minerals were especially more susceptible to reduction compared to their larger counterparts or when subjected to higher applied E_H^{MER} conditions. For instance, Gt_2000 was reduced by 61.7% at -0.35 V, while Gt_30 was reduced completely (100.2%). This increased susceptibility is likely due to the higher redox potential of the smaller particles.

In terms of reduction rate, we chose to present corrected observed rate constants,^{28,43} (k_{obs}^*), for goethite and hematite at E_H^{MER} of -0.35 and -0.53 V, at which all minerals can be reduced to comparatively high extents (from 35.2 to 100.2%). We found that k_{obs}^* values clearly increased with decreasing particle sizes irrespective of E_H^{MER} (Table 1). This demonstrated that the reduction rates were strongly dependent on the mineral particle size. In addition, we calculated the $k_{\text{obs,av}}^{\text{med}}$ for different mediator species used in MER experiments ($k_{\text{obs,av}}^{\text{med}} = 11.19 \pm 0.52$ h⁻¹ for DQ at -0.35 V and 7.75 ± 0.26 h⁻¹ for ZiV at -0.53 V). At $E_H^{\text{MER}} = -0.53$ V, the k_{obs}^* for Gt_30 and Hm_8 reduction (8.61 and 8.00 h⁻¹, respectively; Table 1) was slightly higher than the $k_{\text{obs,av}}^{\text{med}}$ (7.75 h⁻¹). This indicated that the rates of mediator rereduction masked the rates of electron transfer from the reduced mediator to those iron minerals. Consequently, the observed reduction rate constants for Gt_30 and Hm_8 likely do not represent their maximum reduction rate, which could be higher in a case where the mediator re-reduction was faster. Conversely, the rest of the reduction rate constants for the (oxyhydr)oxides were lower than the $k_{\text{obs,av}}^{\text{med}}$, which denoted that the observed reduction

Table 1. Reduction Rate Constants of Goethite and Hematite as a Function of the Particle Size

mineral	microbial reduction ^a h ⁻¹ (×10 ⁻³)	mediated electrochemical reduction	
		$E_{\text{H}}^{\text{MER}} = -0.35 \text{ V}^b$ h ⁻¹	$E_{\text{H}}^{\text{MER}} = -0.53 \text{ V}^c$ h ⁻¹
Gt_2000	1.45 ± 0.03	2.89 ± 0.26	3.95
Gt_90	2.21 ± 0.19	3.38 ± 0.23	4.67
Gt_30	6.05 ± 0.17	8.19 ± 0.16	8.61
Hm_300	0.48 ± 0.02	1.60 ± 0.09	2.20
Hm_40	1.74 ± 0.10	3.55 ± 0.08	4.93
Hm_8	6.20 ± 0.36	7.63 ± 0.31	8.00

^aMicrobial reduction was carried out in triplicates. Values represent the mean ± standard deviation for a given mineral. ^bFor all iron (oxyhydr)oxides at $E_{\text{H}}^{\text{MER}} = -0.35 \text{ V}$, duplicate MER experiments were carried out simultaneously in two separate electrochemical cells. Reported rate constants are the average of the duplicate measurements (± depict deviation of single measurements from the mean). ^cFor all iron (oxyhydr)oxides at $E_{\text{H}}^{\text{MER}} = -0.53 \text{ V}$, MER experiments were performed in individual cells.

rates were controlled by the rates of electron transfer from the reduced mediator to the minerals instead of the redox cycling of the mediators.

In the case of electrochemical reduction, we could quantitatively compare the relative thermodynamic driving force. We first calculated theoretical E_{H} values, which ranged from +765 to +832 mV (for goethite) and from +764 mV to +863 mV (for hematite) (eqs 5 and 6, SI, Table S4).^{53–55} For Hm_40 and Hm_8, the E_{H} values (+781 mV, +863 mV) were similar to previously reported values for similarly sized particles (+773 ± 6 and +823 ± 6 mV).⁵⁸ We then calculated $\Delta_r G$, which yielded values from −49.3 to −60.0 kJ/mol for all goethite and hematite at $E_{\text{H}}^{\text{MER}} = -0.35 \text{ V}$ and from −55.2 to −66.5 kJ/mol at $E_{\text{H}}^{\text{MER}} = -0.53 \text{ V}$ (SI, Section S4). At more negative $\Delta_r G$ values and more positive E_{H} values, the reduction of iron (oxyhydr)oxide was much more thermodynamically favorable, thus leading to a faster and more extensive reduction.

Microbial Fe(III) Reduction of Goethite and Hematite.

To evaluate size-dependent susceptibility of goethite and hematite toward microbial Fe(III) reduction, we tracked the production of Fe(II) over time (Figure 2). With decreasing particle sizes, reduction extents varied from 6.3 to 16.8% for goethite and from 2.7 to 21.7% for hematite (SI, Table S2). The results indicate that smaller-sized minerals were more susceptible toward microbial Fe(III) reduction, which is consistent with mediated electrochemical reduction.

Rate constants (k') for microbial Fe(III) reduction were derived using pseudo-first-order kinetic model (Figure 2), which provided a highly reliable fit to the experimental data ($R^2 > 0.99$ for both samples). Rate constants increased with decreasing particle sizes for both goethite and hematite (Table 1), which is consistent with the trend observed in the mediated electrochemical experiments. Microbial reduction rates were much more sensitive to size compared with mediated electrochemical reduction. For example, Gt_30 was reduced 4.2× faster than Gt_2000 via microbial reduction, compared to only 2.2–2.8× increase for MER (at $E_{\text{H}}^{\text{MER}} = -0.35$ or -0.53 V). The same is true and more pronounced when comparing Hm_8 and Hm_300 (12.9× for microbial and only 3.6–4.8× for electrochemical reduction). To test whether this observed trend is influenced by the presence of AQDS as an electron




Figure 2. Total Fe(II) production by microbial Fe(III) reduction of goethite (blue) (A) and hematite (red) (B) as a function of particle size. Dashed lines denote the fit based on pseudo-first order kinetics. Symbols correspond to the average of biological triplicates, and the error bars indicate standard deviations. Experimental conditions: 10 mM Fe(III) of goethite/hematite; 5×10^8 cells mL⁻¹ *S. oneidensis* MR-1 (no cells in abiotic controls); 20 μM AQDS; pH 7.

shuttle, we also performed microbial experiments in the absence of AQDS. We observed a faster and more extensive reduction of smaller particles, although overall slower than that in the presence of AQDS (Figure S11, Table S3). These findings indicate that microbial reduction is more strongly influenced by the mineral size than the presence of electron shuttles.

Determining Impacts of Related Variables on Mineral Reducibility via MER and Microbial Reduction. The extents and rates of Fe(III) mineral reduction have been suggested to be influenced by a variety of related factors,^{14,18–23} making it challenging to evaluate these variables independently. For example, crystallite size influences both the surface areas and the thermodynamics (i.e., $\Delta_r G$ and E_{H} of the mineral).⁵⁴ Crystallite size and particle concentration also affect the degree and type of aggregation (e.g., pore size), which then further affects the reactive surface areas.^{59,60} In addition, different techniques are able to measure different types of sizes⁶¹ (e.g., crystallite size via XRD, particle size via electron microscopy, and aggregation size via DLS) and surface areas⁶² (e.g., measured via BET or calculated from shapes observed under electron microscopy). Among these variables, surface area and thermodynamics have been strongly proposed to control mineral reduction processes.^{20,28,29} To probe the individual and combined impact of these two variables, we




Figure 3. Results of redundancy analysis (A, C) and Pearson correlation analysis (B, D) for related variables and corresponding reduction extents and rate constants in the mediated electrochemical reduction (A, B) and microbial reduction (C, D). Blue solid lines with solid arrowheads in (A, C) represent reduction extents and rate constants as response variables. Red solid lines with unfilled arrowheads represent different interrelated explanatory variables. Stars in (B, D) represent significant correlation ($P < 0.05$; 0.01 ; 0.001), and the value beside the column represents the extent of correlation.

therefore performed a correlation analysis between rate constant normalized using BET-derived surface area and free reaction energy ($\Delta_r G$) or standard redox potential (E_H^0) (SI, Figure S12). Our results indicated that neither variable alone nor their combination can fully account for the observed variations in reduction rate constants. Thus, a data analysis technique that incorporates these related variables (also called “explanatory variables”) is necessary.

To clarify the impact of mineral-related variables on their reduction extents and rates, redundancy analysis (RDA) was applied to the data collected from mediated electrochemical and microbial reduction experiments (SI, Section S4, Tables S4 and S5). To complement RDA, the significance of correlation was also analyzed through Pearson correlation analysis.⁶³

For mediated electrochemical reduction, the greatest impacts on rate and extent reflected by RDA were $\Delta_r G$, E_H , TEM surface area, BET surface area, and crystallite size (Figure 3A). The acute angles ($<90^\circ$) between red arrows for E_H , BET surface area, TEM surface area, surface charge, and particle number with rate and extent reflected their positive correlations, and the obtuse angles ($>90^\circ$) between red arrows for $\Delta_r G$ and crystallite size with rate and extent indicated their negative correlations. The significance of correlations between the above interrelated variables and rate and extent is shown in Figure 3B ($P < 0.05$, 0.01 , 0.005). The six variables mentioned

above (particle number, crystallite size, $\Delta_r G$, TEM SA, BET SA, and E_H) were all significantly correlated to both reduction rate and extent. In comparison, for microbial reduction experiments, the greatest impacts were E_H , crystallite size, reactive sites, and BET SA and TEM SA (Figure 3C). Significant correlations were only found between rate and extent and particle number, crystallite size, reactive sites, aggregation size, and E_H (Figure 3D).

Collectively, the results from RDA and Pearson correlation analysis indicated that the thermodynamic driving force (represented by $\Delta_r G$ and/or E_H) and surface processes (represented by TEM surface areas (for MER) and reactive sites (for microbial Fe(III) reduction)) are the main controlling factors in reduction. This goes beyond the current understanding as past work has shown either (a) BET-based surface area is the only controlling factor^{18,20} or (b) when comparing different bulk minerals that thermodynamic driving force is the most important.^{28,29} Here, we show that thermodynamics plays a primary role, while surface area plays a secondary role, which can be applied to varying particle size and different mineral types.

Our results are supported by previous studies that highlighted thermodynamics and surface availability in controlling Fe(III) reduction. From a theoretical thermodynamics viewpoint, small particles are expected to have higher free energy and reduction potential than their large counter-




Figure 4. Reduced iron (Fe(II)) production was based on the isotopic composition of the microbially reduced Fe from reactors containing an isotopically labeled mix of different sizes of goethite and hematite. Data points correspond to an average of biological triplicates, and the error bars indicate standard deviations. Pie chart inside each panel illustrates fractions of Fe(II) derived from specific sized minerals at the final time point. Experimental conditions: 5×10^8 cells mL^{-1} *S. oneidensis* MR-1; 20 μM AQDS; pH 7; (A) 5 mM $^{56}\text{Gt}_{2000}$ and 5 mM $^{\text{NA}}\text{Gt}_{90}$; (B) 3.5 mM $^{56}\text{Hm}_{40}$ and 6.5 mM $^{\text{NA}}\text{Hm}_8$; (C) 4.5 mM $^{\text{NA}}\text{Gt}_{90}$ and 5.5 mM $^{56}\text{Hm}_8$.

parts,^{54,55} thus making them more reactive and allowing for faster rates and higher extent of Fe(III) oxide reduction. This hypothesis was supported by the relationship between the reduction behavior and our calculated E_{H} and $\Delta_r G$ values.

Additionally, surface area has been identified as a key controlling factor for both biotic and abiotic Fe(III) reduction in several papers.^{18,20} Our MER results also showed a positive correlation between surface area and Fe(III) reduction rates. Interestingly, in our microbial Fe(III) reduction experiments, it was the reactive site rather than the BET or TEM surface area that contributed more to the reduction rate and extent. Past studies^{24,64} have suggested that nanoparticles are prone to aggregation, strongly affecting their reactive surface area and further reactivity. Our batch MER experiments were carried out at a low initial Fe(III) concentration and with continuous stirring. Therefore, it is possible that goethite and hematite particles underwent slight aggregation, and thus, the reactive surface area may still be reasonably represented by the TEM and BET surface area measurements. Conversely, the significantly higher initial Fe(III) concentration (70× higher relative to MER experiments) and the stationary conditions maintained throughout the experiments can promote extensive nanoparticle aggregation. Further aggregation can be promoted by the presence of microbial cells and potential extracellular polymeric substances (EPS), which was supported by the tight association between cells and small particle aggregates (Gt₉₀ and Hm₈) and relatively loose association with large particle aggregates (Gt₂₀₀₀ and Hm₄₀) determined by fluorescence microscopy (Figure S14). This aggregation may decrease the reactive sites on the nanoparticle surfaces, thereby limiting the accessibility of AQDS to these sites. Herein, the adsorption of Fe(II) by particles was measured and used as a proxy for reactive sites, so that the effect of aggregation on reactive sites could be accounted for. The RDA results demonstrated reactive sites correlated better with microbial Fe(III) reduction, compared to surface area.

When applying these results to the natural environment, note that environmental factors such as temperature, pH, and microbial community may affect reduction rate constants.^{11,12,65} For instance, relatively lower pH may promote reduction due to the availability of protons and more exposed reactive sites.⁶⁶

Microbial Fe(III) Reduction in Mixed Mineral Systems. An experimental system with iron (oxyhydr)oxides of different

isotopic compositions allowed us to track the origin of the Fe reduction during incubation with microorganisms. In mixed mineral systems, we first performed linear combination between $E_{\text{H,mix}}$ and SA_{mix} and overall rate constants, which suggest no clear tendency compared to single mineral systems (SI, Figure S15). To determine whether smaller particles of goethite and hematite are preferentially reduced by *S. oneidensis* MR-1 in mixed mineral systems, we calculated the fraction of Fe reduced from specific sized minerals based on the ^{54}Fe concentration of the total Fe(II) produced (details in SI, Section S5). We observed that although both minerals in each mixed mineral system were reduced simultaneously, the majority of the Fe(II) produced, ranging from 81 to 97%, was attributed to the reduction of smaller minerals (Figure 4). We further determined both reduction extents and rate constants for each mineral to compare between single and mixed mineral systems (SI). Table S6 shows that the reduction extents of small minerals are always higher than those of large minerals, which is overall consistent with the previous single mineral systems (Tables 1 and S2). In addition, the reduction extents of small minerals in mixed mineral systems are slightly higher than their reduction extent in single mineral systems under the same environmental conditions. For instance, Hm₈ in Hm/Gt mixed systems is reduced 2× more compared to single mineral systems. In contrast, the reduction extents of large minerals are much lower than those in single mineral systems. Specifically, Gt₂₀₀₀, Hm₄₀ and Gt₉₀ in mixed mineral systems are reduced $3 \times$, $6 \times$ and $5 \times$ less than in single mineral systems. Figure S16 shows the fraction of Fe(II) derived from each mineral in mixed mineral systems, demonstrating that 81, 97, and 97% of Fe(II) can be attributed to reduction of smaller minerals. These findings confirmed that smaller minerals are preferentially reduced by *S. oneidensis* MR-1 in mixed mineral systems and even to a higher extent than that solely predicted based on single mineral systems.

The different reduction behaviors can also be observed in the reduction rate constants (SI, Table S6). When considering single mineral systems, reduction rate constants of Gt₉₀, Hm₈, and Hm₈ were 1.5×, 3.6×, and 3.2× greater than those of Gt₂₀₀₀, Hm₄₀, and Gt₉₀, respectively. When reduction happened in mixed mineral systems, reduction rate constants of Gt₉₀, Hm₈, and Hm₈ were 6×, 21×, and 23× greater than their large-sized counterparts. This significant difference was attributed mostly to a pronounced reduction

suppression of large particles and secondarily to a slight promotion of reduction of small particles.

Possible Mechanisms for Preferential Microbial Reduction of Small-Sized Particles over Large Particles in Mixed Mineral Systems. Based on our RDA and Pearson correlation analysis above, E_H and reactive sites were the most important variables for microbial reduction in single mineral systems. Thus, they also likely contribute to differences in the reduction extent and rate in our mixed mineral systems. In our mixed mineral systems, the reduction of small particles is expected to be more thermodynamically favorable (higher E_H values). In addition, given the initially added concentration and surface area of specific iron (oxyhydr)oxides in mixed systems, small particles had 2× to 7× higher reactive sites compared to large particles, which can contribute to differential reduction.

The potential preferential adsorption of AQDS between particles may also contribute to differential reduction of the Fe(III) mineral. Negatively charged AQDS acts as an electron shuttle between Fe(III)-reducing bacteria and minerals.^{66,67} The point of zero charge of our synthesized minerals were 9.2 and 8.5 for Gt_2000 and Gt_90, and 9.2 and 8.0 for Hm_40 and Hm_8, respectively (SI, Figure S17). It has been extensively shown that negatively charged organic molecules adsorb on iron (oxyhydr)oxide via electrostatic interactions.^{68,69} Thus, small particles which exhibit more positively charged surfaces than their large counterparts at our experimental pH (7.0) may provide higher electrostatic attraction for AQDS. In addition, the number of reactive surface sites ($>10^{18}$) on small particles is an order of magnitude higher than that of added AQDS molecules (10^{17}) (SI, Table S5). If each reactive site can bind only one AQDS molecule, then there will be insufficient AQDS to react with large particles. Consequently, this could inhibit the reduction of large particles in mixed mineral systems.

The presence of defects may also lead to a preferential reduction. It has been found that defects can facilitate electron transfer in both abiotic and biotic systems.^{38,70} In our study, the presence of defects in small particles was confirmed through XRD and Mössbauer spectroscopy data. These defects likely contribute to the increased electron transfer in smaller particles, making them more prone to reduction.

Possible Mechanisms for Suppressed Microbial Reduction of Large-Sized Particles in Mixed Mineral Systems Compared to Single Mineral Systems. Interestingly, the observed pronounced suppression of reduction in large-sized particles occurs not only in comparison to small particles but also when compared to particles of the same size within homogeneous single mineral systems. In a system with a broad particle size distribution, such as the mixed mineral system in this study, Ostwald ripening is a process worth considering. As far as we could measure, Ostwald ripening did not play a major role. We found no clear evidence that large particles in mixed mineral systems became noticeably larger (Figure S18), when compared to those in single mineral systems. This is supported by our earlier findings^{71,72} that the length of the particles of nanoparticulate goethite does not change substantially, even in the presence of Fe^{2+} . Below, we propose three explanations for suppressed reduction of large particles (SI, Figure S20).

Herein, we speculate that preferential reduction can be ascribed to varied aggregation patterns in mixed mineral systems. Mineral mixtures can result in heteroaggregation due to their morphological variations and surface charge disequi-

librium.^{41,73,74} In our present study, the mixed minerals displayed shifts toward larger aggregate sizes relative to the single minerals (Detailed data and description in SI, Section S6). This could result in increased physical coverage of reactive sites and a decrease in the aggregate pore throat, which slows diffusion of shuttle molecules such as AQDS (0.5–1.5 nm size) (SI, Section S6). However, it must be noted that DLS data can indicate only the hydrodynamic diameter of an aggregate in the absence of microbes, without any knowledge on the aggregate morphology and compaction factor (i.e., loose versus compact packing) as well as its association with microbial cells. Further exploration using cryogenic transmission electron microscopy (cryo-TEM) and small-angle X-ray scattering (SAXS) is needed to test this hypothesis.

The second speculation is related to the electron transfer (ET) process, leading to the impeded reduction of large particles and the enhanced reduction of small particles. Previous studies have substantiated the existence of electron transfer via electron hopping between different microbial species,^{75–78} between minerals and minerals,^{79,80} as well as within iron (oxyhydr)oxides.⁸¹ More recently, it has been revealed that directional long-distance ET can occur along a redox gradient, driven by differences in reduction potentials, with electrons moving from lower to higher reduction potentials.⁸² Therefore, large particles with lower reduction potentials can facilitate directional electron transfer toward small ones, when these minerals are present as heteroaggregates.

The third speculation is attributed to the disproportionate adsorption capacity and affinity toward dissolved Fe(II) among different sized minerals. It has been reported that dissolved Fe(II) adsorbs onto residual minerals, thereby inhibiting their further reduction.^{19,83} As discussed above, large particles were less positively charged than small ones, as indicated by the zeta potentials of 29.8 mV for Hm_40 and 34.5 mV for Hm_8 (SI, Table S5). The less positive surface charge can lead to released Fe^{2+} being more easily attracted to their surface. This selective attraction may block reactive surface sites, thereby forming a passivated layer that inhibits further reduction. This passivation effect may result in slower and less extensive reduction of large particles in mixed mineral systems, compared to single mineral systems.

ENVIRONMENTAL IMPLICATIONS

The reduction of iron (oxyhydr)oxides is one of the most important biogeochemical processes on Earth that significantly impacts the fate and bioavailability of nutrients, trace elements and pollutants.^{7,12} Iron(III) mineral reduction was originally thought to be predominantly influenced by mineral phases and crystallinity. Poorly crystalline minerals, like ferrihydrite, were found to be much more susceptible to reduction compared to highly crystalline minerals, such as lepidocrocite, magnetite, goethite and hematite.^{14,21,24,84,85} Our results show that particle size plays an important role in controlling reduction kinetics, potentially outweighing other factors such as crystallinity and the mineral phase. For instance, hematite (8 nm) in our studies was more susceptible to reduction compared to goethite of larger sizes (30, 90, and 2000 nm), regardless of electrochemical and microbial reduction. In addition, in heterogeneous environments where minerals of varying sizes and phases are likely to be present together, our results suggested an enhancement of the reduction of small

particles and a strong suppression of the reduction of larger particles.

Our findings have significant implications for the fate and behavior of nutrients, pollutants, and trace elements associated with iron (oxyhydr)oxide minerals. Iron (oxyhydr)oxides have been widely employed in environmental remediation and agricultural production to remove contaminants and enhance soil fertility, due to their high sorption capacities.^{86–88} However, nanoparticles are more susceptible to undergoing extensive reduction under anoxic conditions, affecting the mobility and bioavailability of trace elements and pollutants in soils and aquatic systems. These interactions suggest that when accounting for particle size of iron minerals, more accurate predictions of trace element behavior under redox fluctuating conditions can be made. In addition, the higher reactivity of smaller particles should be incorporated into reactive transport models to better estimate the flux of elements across the redox boundaries in natural systems.

Environmental studies focused on particles suspended in aquatic bodies commonly utilize 0.45 or 0.22 μm membrane filters to separate solid and dissolved aqueous phases, which fails to account for nanoparticles that can pass through filters. To illustrate the impact of particle size on the reducible phase, we developed a model assuming a total Fe concentration of 10 mM, with particle sizes distributed across three ranges: 1–10, 10–100, and >100 nm. By applying reduction extents from our experiments, we approximated the reducible fraction for each size range, allowing us to calculate the contributions of different particle sizes to the overall reducible Fe phase (details in SI, Section S7). The model results clearly demonstrate that neglecting the size effect and not considering particles <0.45 μm can lead to a significant underestimation (7 $\times$ for hematite and 14 $\times$ for goethite) of the reducible solid phase in natural aquatic systems (Figure S21). This highlights the need for more advanced techniques, such as field-flow fractionation (FFF) and single particle inductively coupled plasma mass spectrometry (SP-ICP-MS), which can more accurately detect and characterize nanoparticles. Furthermore, it is important to recognize that naturally formed iron (oxyhydr)oxides may exhibit different behaviors than synthesized iron oxides under laboratory conditions. Therefore, future field studies are required to validate the behavior of natural iron (oxyhydr)oxides under complex environmental conditions.

■ ASSOCIATED CONTENT

SI Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.est.4c08032>.

Details about synthesis and characterization of nanoparticulate goethite and hematite, mediated electrochemical reduction of iron oxides, data collection and calculation for RDA and Pearson correlation analysis, and calculation of the origin of reduced Fe based on their isotope concentration (PDF)

■ AUTHOR INFORMATION

Corresponding Author

Prachi Joshi – Geomicrobiology, Department of Geosciences, University of Tübingen, 72076 Tübingen, Germany; orcid.org/0000-0001-5954-0309; Email: Prachi.joshi@uni-tuebingen.de

Authors

Xiyang Xu – Geomicrobiology, Department of Geosciences, University of Tübingen, 72076 Tübingen, Germany

Muammar Mansor – Geomicrobiology, Department of Geosciences, University of Tübingen, 72076 Tübingen, Germany; orcid.org/0000-0001-7830-650X

Guoxiang Li – Environmental Chemistry and Mineralogy, Department of Geosciences, University of Tübingen, 72076 Tübingen, Germany

Tsz Ho Chiu – Geomicrobiology, Department of Geosciences, University of Tübingen, 72076 Tübingen, Germany; orcid.org/0000-0003-4734-6490

Stefan B. Haderlein – Environmental Chemistry and Mineralogy, Department of Geosciences, University of Tübingen, 72076 Tübingen, Germany

Andreas Kappler – Geomicrobiology, Department of Geosciences, University of Tübingen, 72076 Tübingen, Germany; Cluster of Excellence: EXC 2124: Controlling Microbes to Fight Infection, 72076 Tübingen, Germany; orcid.org/0000-0002-3558-9500

Complete contact information is available at:

<https://pubs.acs.org/doi/10.1021/acs.est.4c08032>

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We thank Franziska Schaedler and Lars Grimm for their technical support in the lab, Carolin Dreher for her XRD measurements, and Martina Bottaro for her ICP-MS analysis. We are grateful to Jeremiah Shuster for his assistance with SEM measurements, to Sandra Richter, Rebecca Stahl, and Lorenz Henneberg for their assistance with TEM measurements, also to the Tübingen Structural Microscopy Core Facility (TSM) and Center for Molecular Biology of Plants (ZMBP) for their support (DFG INST 37/900-1 FUGG). We are grateful for infrastructural support by the DFG under Germany's Excellence Strategy, cluster of Excellence EXC2124 (project ID 390838134). This research was supported by the CSC scholarship (202106010075). We would also like to thank the reviewers whose input greatly improved the manuscript.

■ REFERENCES

- (1) Cornell, R. M.; Schwertmann, U. *The iron oxides: structure, properties, reactions, occurrences, and uses*; Wiley-Vch Weinheim, 2003; Vol. 664.
- (2) Hochella, M. F.; Lower, S. K.; Maurice, P. A.; Penn, R. L.; Sahai, N.; Sparks, D. L.; Twining, B. S. Nanominerals, mineral nanoparticles, and earth systems. *Science* **2008**, 319 (5870), 1631–1635.
- (3) Hochella, M. F.; Mogk, D. W.; Ranville, J.; Allen, I. C.; Luther, G. W.; Marr, L. C.; McGrail, B. P.; Murayama, M.; Qafoku, N. P.; Rosso, K. M. Natural, incidental, and engineered nanomaterials and their impacts on the Earth system. *Science* **2019**, 363 (6434), No. eaau8299.
- (4) Banfield, J. F.; Zhang, H. Nanoparticles in the environment. *Rev. Mineral. Geochem.* **2001**, 44 (1), 1–58.
- (5) Stumm, W.; Sulzberger, B. The cycling of iron in natural environments: considerations based on laboratory studies of heterogeneous redox processes. *Geochim. Cosmochim. Acta* **1992**, 56 (8), 3233–3257.
- (6) Jickells, T.; An, Z.; Andersen, K. K.; Baker, A.; Bergametti, G.; Brooks, N.; Cao, J.; Boyd, P.; Duce, R.; Hunter, K. Global iron

connections between desert dust, ocean biogeochemistry, and climate. *Science* **2005**, *308* (5718), 67–71.

(7) Borch, T.; Kretzschmar, R.; Kappler, A.; Cappellen, P. V.; Ginder-Vogel, M.; Voegelin, A.; Campbell, K. Biogeochemical redox processes and their impact on contaminant dynamics. *Environ. Sci. Technol.* **2010**, *44* (1), 15–23.

(8) Raiswell, R.; Canfield, D. E. The iron biogeochemical cycle past and present. *Geochim. Perspect.* **2012**, *1* (1), 1–2.

(9) Colombo, C.; Palumbo, G.; He, J.-Z.; Pinton, R.; Cesco, S. Review on iron availability in soil: interaction of Fe minerals, plants, and microbes. *J. Soils Sediments* **2014**, *14*, 538–548.

(10) Melton, E. D.; Swanner, E. D.; Behrens, S.; Schmidt, C.; Kappler, A. The interplay of microbially mediated and abiotic reactions in the biogeochemical Fe cycle. *Nat. Rev. Microbiol.* **2014**, *12* (12), 797–808.

(11) Kappler, A.; Bryce, C.; Mansor, M.; Lueder, U.; Byrne, J. M.; Swanner, E. D. An evolving view on biogeochemical cycling of iron. *Nat. Rev. Microbiol.* **2021**, *19* (6), 360–374.

(12) Weber, K. A.; Achenbach, L. A.; Coates, J. D. Microorganisms pumping iron: anaerobic microbial iron oxidation and reduction. *Nat. Rev. Microbiol.* **2006**, *4* (10), 752–764.

(13) Richter, K.; Schicklberger, M.; Gescher, J. Dissimilatory reduction of extracellular electron acceptors in anaerobic respiration. *Appl. Environ. Microbiol.* **2012**, *78* (4), 913–921.

(14) Cutting, R.; Coker, V.; Fellowes, J.; Lloyd, J.; Vaughan, D. Mineralogical and morphological constraints on the reduction of Fe (III) minerals by *Geobacter sulfurreducens*. *Geochim. Cosmochim. Acta* **2009**, *73* (14), 4004–4022.

(15) Konhauser, K. O.; Kappler, A.; Roden, E. E. Iron in microbial metabolisms. *Elements* **2011**, *7* (2), 89–93.

(16) Pedersen, H. D.; Postma, D.; Jakobsen, R. Release of arsenic associated with the reduction and transformation of iron oxides. *Geochim. Cosmochim. Acta* **2006**, *70* (16), 4116–4129.

(17) Frierdich, A. J.; Catalano, J. G. Controls on Fe (II)-activated trace element release from goethite and hematite. *Environ. Sci. Technol.* **2012**, *46* (3), 1519–1526.

(18) Roden, E. E.; Zachara, J. M. Microbial reduction of crystalline iron (III) oxides: influence of oxide surface area and potential for cell growth. *Environ. Sci. Technol.* **1996**, *30* (5), 1618–1628.

(19) Liu, C.; Zachara, J. M.; Gorby, Y. A.; Szecsody, J. E.; Brown, C. F. Microbial reduction of Fe (III) and sorption/precipitation of Fe (II) on *Shewanella putrefaciens* strain CN32. *Environ. Sci. Technol.* **2001**, *35* (7), 1385–1393.

(20) Roden, E. E. Fe (III) oxide reactivity toward biological versus chemical reduction. *Environ. Sci. Technol.* **2003**, *37* (7), 1319–1324.

(21) O'Loughlin, E. J. Effects of electron transfer mediators on the bioreduction of lepidocrocite (γ -FeOOH) by *Shewanella putrefaciens* CN32. *Environ. Sci. Technol.* **2008**, *42* (18), 6876–6882.

(22) Echigo, T.; Aruguete, D. M.; Murayama, M.; Hochella, M. F., Jr. Influence of size, morphology, surface structure, and aggregation state on reductive dissolution of hematite nanoparticles with ascorbic acid. *Geochim. Cosmochim. Acta* **2012**, *90*, 149–162.

(23) Liu, J.; Pearce, C. I.; Shi, L.; Wang, Z.; Shi, Z.; Arenholz, E.; Rosso, K. M. Particle size effect and the mechanism of hematite reduction by the outer membrane cytochrome OmcA of *Shewanella oneidensis* MR-1. *Geochim. Cosmochim. Acta* **2016**, *193*, 160–175.

(24) Bose, S.; Hochella, M. F., Jr.; Gorby, Y. A.; Kennedy, D. W.; McCready, D. E.; Madden, A. S.; Lower, B. H. Bioreduction of hematite nanoparticles by the dissimilatory iron reducing bacterium *Shewanella oneidensis* MR-1. *Geochim. Cosmochim. Acta* **2009**, *73* (4), 962–976.

(25) Bonneville, S.; Van Cappellen, P.; Behrends, T. Microbial reduction of iron (III) oxyhydroxides: effects of mineral solubility and availability. *Chem. Geol.* **2004**, *212* (3–4), 255–268.

(26) Bonneville, S.; Behrends, T.; Van Cappellen, P.; Hyacinthe, C.; Röling, W. F. Reduction of Fe (III) colloids by *Shewanella putrefaciens*: a kinetic model. *Geochim. Cosmochim. Acta* **2006**, *70* (23), 5842–5854.

(27) Yan, B.; Wrenn, B. A.; Basak, S.; Biswas, P.; Giammar, D. E. Microbial reduction of Fe (III) in hematite nanoparticles by *Geobacter sulfurreducens*. *Environ. Sci. Technol.* **2008**, *42* (17), 6526–6531.

(28) Aeppli, M.; Voegelin, A.; Gorski, C. A.; Hofstetter, T. B.; Sander, M. Mediated electrochemical reduction of iron (oxyhydr-) oxides under defined thermodynamic boundary conditions. *Environ. Sci. Technol.* **2018**, *52* (2), 560–570.

(29) Aeppli, M.; Giroud, S.; Vranic, S.; Voegelin, A.; Hofstetter, T. B.; Sander, M. Thermodynamic controls on rates of iron oxide reduction by extracellular electron shuttles. *Proc. Natl. Acad. Sci. U. S. A.* **2022**, *119* (3), No. e2115629119.

(30) Schwertmann, U. Solubility and dissolution of iron oxides. *Plant Soil* **1991**, *130*, 1–25.

(31) Regelin, I. C.; Voegelin, A.; Weng, L.; Koopmans, G. F.; Comans, R. N. Characterization of colloidal Fe from soils using field-flow fractionation and Fe K-edge X-ray absorption spectroscopy. *Environ. Sci. Technol.* **2014**, *48* (8), 4307–4316.

(32) Canfield, D. E. Reactive iron in marine sediments. *Geochim. Cosmochim. Acta* **1989**, *53* (3), 619–632.

(33) van der Zee, C.; Roberts, D. R.; Rancourt, D. G.; Slomp, C. P. Nanogoethite is the dominant reactive oxyhydroxide phase in lake and marine sediments. *Geology* **2003**, *31* (11), 993–996.

(34) Balsam, W.; Ji, J.; Chen, J. Climatic interpretation of the Luochuan and Lingtai loess sections, China, based on changing iron oxide mineralogy and magnetic susceptibility. *Earth Planet. Sci. Lett.* **2004**, *223* (3–4), 335–348.

(35) Scheinost, A.; Chavernas, A.; Barrón, V.; Torrent, J. Use and limitations of second-derivative diffuse reflectance spectroscopy in the visible to near-infrared range to identify and quantify Fe oxide minerals in soils. *Clay Clay Min.* **1998**, *46* (5), 528–536.

(36) Ranville, J.; Montano, M. D., Size distributions. In *Frontiers of Nanoscience*; Elsevier, 2015; Vol. 8, pp 91–121.

(37) Westerhoff, P.; Atkinson, A.; Fortner, J.; Wong, M. S.; Zimmerman, J.; Gardea-Torresdey, J.; Ranville, J.; Herckes, P. Low risk posed by engineered and incidental nanoparticles in drinking water. *Nat. Nanotechnol.* **2018**, *13* (8), 661–669.

(38) Notini, L.; Byrne, J. M.; Tomaszewski, E. J.; Latta, D. E.; Zhou, Z.; Scherer, M. M.; Kappler, A. Mineral defects enhance bioavailability of goethite toward microbial Fe (III) reduction. *Environ. Sci. Technol.* **2019**, *53* (15), 8883–8891.

(39) Hua, J.; Feng, C.; Sun, J.; Wu, F.; Wu, F.; Liu, C. Facet-preferential reduction of hematite nanocrystals by *Shewanella oneidensis* MR-1: An iron isotope tracer study. *Chem. Geol.* **2022**, *614*, No. 121166.

(40) Notini, L.; ThomasArrigo, L. K.; Kaegi, R.; Kretzschmar, R. Coexisting goethite promotes Fe (II)-catalyzed transformation of ferrihydrite to goethite. *Environ. Sci. Technol.* **2022**, *56* (17), 12723–12733.

(41) Ma, J.; Jing, Y.; Gao, L.; Chen, J.; Wang, Z.; Weng, L.; Li, H.; Chen, Y.; Li, Y. Hetero-aggregation of goethite and ferrihydrite nanoparticles controlled by goethite nanoparticles with elongated morphology. *Sci. Total Environ.* **2020**, *748*, No. 141536.

(42) Gorski, C. A.; Edwards, R.; Sander, M.; Hofstetter, T. B.; Stewart, S. M. Thermodynamic characterization of iron oxide–aqueous Fe²⁺ redox couples. *Environ. Sci. Technol.* **2016**, *50* (16), 8538–8547.

(43) Sander, M.; Hofstetter, T. B.; Gorski, C. A. Electrochemical analyses of redox-active iron minerals: a review of nonmediated and mediated approaches. *Environ. Sci. Technol.* **2015**, *49* (10), 5862–5878.

(44) Anschutz, A. J.; Penn, R. L. Reduction of crystalline iron (III) oxyhydroxides using hydroquinone: Influence of phase and particle size. *Geochim. Trans.* **2005**, *6* (3), 60.

(45) Madden, A. S.; Hochella, M. F., Jr.; Luxton, T. P. Insights for size-dependent reactivity of hematite nanomineral surfaces through Cu²⁺ sorption. *Geochim. Cosmochim. Acta* **2006**, *70* (16), 4095–4104.

(46) Schwertmann, U.; Cornell, R. M. *Iron oxides in the laboratory: preparation and characterization*; John Wiley & Sons, 2008.

- (47) Lowry, G. V.; Hill, R. J.; Harper, S.; Rawle, A. F.; Hendren, C. O.; Klaessig, F.; Nobbmann, U.; Sayre, P.; Rumble, J. Guidance to improve the scientific value of zeta-potential measurements in nanoEHS. *Environ. Sci. Nano* **2016**, 3 (5), 953–965.
- (48) Fultz, M. L.; Durst, R. A. Mediator compounds for the electrochemical study of biological redox systems: a compilation. *Anal. Chim. Acta* **1982**, 140 (1), 1–18.
- (49) Gorski, C. A.; Klüpfel, L.; Voegelin, A.; Sander, M.; Hofstetter, T. B. Redox properties of structural Fe in clay minerals. 2. Electrochemical and spectroscopic characterization of electron transfer irreversibility in ferruginous smectite, SWa-1. *Environ. Sci. Technol.* **2012**, 46 (17), 9369–9377.
- (50) Cross, S. N.; Al-Feghali, A. H.; Blum, A. S. Interactions of Common Biological Buffers with Iron Oxide Nanoparticles. *Langmuir* **2023**, 39 (22), 7632–7641.
- (51) Stookey, L. L. Ferrozine—a new spectrophotometric reagent for iron. *Anal. Chem.* **1970**, 42 (7), 779–781.
- (52) Hegler, F.; Posth, N. R.; Jiang, J.; Kappler, A. Physiology of phototrophic iron (II)-oxidizing bacteria: implications for modern and ancient environments. *FEMS Microbiology Ecology* **2008**, 66 (2), 250–260.
- (53) Robie, R. A.; Hemingway, B. S. *Thermodynamic properties of minerals and related substances at 298.15 K and 1 bar (105 Pascals) pressure and at higher temperatures*; US Government Printing Office, 1995; Vol. 2131.
- (54) Navrotsky, A.; Mazeina, L.; Majzlan, J. Size-driven structural and thermodynamic complexity in iron oxides. *Science* **2008**, 319 (5870), 1635–1638.
- (55) Navrotsky, A.; Ma, C.; Lilova, K.; Birkner, N. Nanophase transition metal oxides show large thermodynamically driven shifts in oxidation-reduction equilibria. *Science* **2010**, 330 (6001), 199–201.
- (56) Davies, C. W. 397. The extent of dissociation of salts in water. Part VIII. An equation for the mean ionic activity coefficient of an electrolyte in water, and a revision of the dissociation constants of some sulphates. *J. Chem. Soc. (Resumed)* **1938**, 2093–2098.
- (57) Zhang, L.; Yang, H.; Tang, J.; Qiu, X.; Yu, A. Y. Attenuation of arsenic in a karst subterranean stream and correlation with geochemical factors: A case study at Lihu, South China. *J. Environ. Sci.* **2014**, 26 (11), 2222–2230.
- (58) Chen, G.; Thompson, A.; Gorski, C. A. Disentangling the size-dependent redox reactivity of iron oxides using thermodynamic relationships. *Proc. Natl. Acad. Sci. U. S. A.* **2022**, 119 (40), No. e2204673119.
- (59) Jeong, H. Y.; Lee, J. H.; Hayes, K. F. Characterization of synthetic nanocrystalline mackinawite: crystal structure, particle size, and specific surface area. *Geochim. Cosmochim. Acta* **2008**, 72 (2), 493–505.
- (60) Liu, J.; Aruguete, D. M.; Murayama, M.; Hochella, M. F., Jr Influence of size and aggregation on the reactivity of an environmentally and industrially relevant nanomaterial (PbS). *Environ. Sci. Technol.* **2009**, 43 (21), 8178–8183.
- (61) Moens, C.; Waegeneers, N.; Fritzsche, A.; Nobels, P.; Smolders, E. A systematic evaluation of Flow Field Flow Fractionation and single-particle ICP-MS to obtain the size distribution of organomineral iron oxyhydroxide colloids. *J. Chromatogr. A* **2019**, 1599, 203–214.
- (62) Lowell, S.; Shields, J. E.; Thomas, M. A.; Thommes, M. *Characterization of porous solids and powders: surface area, pore size and density*; Springer Science & Business Media, 2012; Vol. 16.
- (63) Sedgwick, P. Pearson's correlation coefficient. *BMJ* **2012**, 345, No. e4483.
- (64) Cwintny, D. M.; Handler, R. M.; Schaefer, M. V.; Grassian, V. H.; Scherer, M. M. Interpreting nanoscale size-effects in aggregated Fe-oxide suspensions: reaction of Fe (II) with goethite. *Geochim. Cosmochim. Acta* **2008**, 72 (5), 1365–1380.
- (65) Jin, Q.; Kirk, M. F. pH as a primary control in environmental microbiology: 1. Thermodynamic perspective. *Front. Environ. Sci.* **2018**, 6, 21.
- (66) Lovley, D. R.; Coates, J. D.; Blunt-Harris, E. L.; Phillips, E. J.; Woodward, J. C. Humic substances as electron acceptors for microbial respiration. *Nature* **1996**, 382 (6590), 445–448.
- (67) Bai, Y.; Mella, A.; Cirpka, O. A.; Sun, T.; Angenent, L. T.; Haderlein, S. B.; Kappler, A. AQDS and redox-active NOM enables microbial Fe (III)-mineral reduction at cm-scales. *Environ. Sci. Technol.* **2020**, 54 (7), 4131–4139.
- (68) Fang, L.; Cao, Y.; Huang, Q.; Walker, S. L.; Cai, P. Reactions between bacterial exopolymers and goethite: A combined macroscopic and spectroscopic investigation. *Water Res.* **2012**, 46 (17), 5613–5620.
- (69) Parikh, S. J.; Chorover, J. ATR-FTIR spectroscopy reveals bond formation during bacterial adhesion to iron oxide. *Langmuir* **2006**, 22 (20), 8492–8500.
- (70) Notini, L.; Latta, D. E.; Neumann, A.; Pearce, C. I.; Sassi, M.; N'Diaye, A. T.; Rosso, K. M.; Scherer, M. M. The role of defects in Fe (II)–goethite electron transfer. *Environ. Sci. Technol.* **2018**, 52 (5), 2751–2759.
- (71) Joshi, P.; Fantle, M. S.; Larese-Casanova, P.; Gorski, C. A. Susceptibility of goethite to Fe²⁺-catalyzed recrystallization over time. *Environ. Sci. Technol.* **2017**, 51 (20), 11681–11691.
- (72) Joshi, P.; Gorski, C. A. Anisotropic morphological changes in goethite during Fe²⁺-catalyzed recrystallization. *Environ. Sci. Technol.* **2016**, 50 (14), 7315–7324.
- (73) Wang, D.; Jin, Y.; Jaisi, D. P. Effect of size-selective retention on the cotransport of hydroxyapatite and goethite nanoparticles in saturated porous media. *Environ. Sci. Technol.* **2015**, 49 (14), 8461–8470.
- (74) Gupta, G. S.; Senapati, V. A.; Dhawan, A.; Shanker, R. Heteroagglomeration of zinc oxide nanoparticles with clay mineral modulates the bioavailability and toxicity of nanoparticle in *Tetrahymena pyriformis*. *J. Colloid Interface Sci.* **2017**, 495, 9–18.
- (75) Kato, S.; Hashimoto, K.; Watanabe, K. Microbial interspecies electron transfer via electric currents through conductive minerals. *Proc. Natl. Acad. Sci. U. S. A.* **2012**, 109 (25), 10042–10046.
- (76) Nakamura, R.; Kai, F.; Okamoto, A.; Newton, G. J.; Hashimoto, K. Self-constructed electrically conductive bacterial networks. *Angew. Chem.-Int. Ed.* **2009**, 121 (3), 516–519.
- (77) Shi, L.; Dong, H.; Reguera, G.; Beyenal, H.; Lu, A.; Liu, J.; Yu, H.-Q.; Fredrickson, J. K. Extracellular electron transfer mechanisms between microorganisms and minerals. *Nat. Rev. Microbiol.* **2016**, 14 (10), 651–662.
- (78) Xu, H.; Chang, J.; Wang, H.; Liu, Y.; Zhang, X.; Liang, P.; Huang, X. Enhancing direct interspecies electron transfer in syntrophic-methanogenic associations with (semi) conductive iron oxides: Effects and mechanisms. *Sci. Total Environ.* **2019**, 695, No. 133876.
- (79) Liao, W.; Ye, Z.; Yuan, S.; Cai, Q.; Tong, M.; Qian, A.; Cheng, D. Effect of coexisting Fe (III)(oxyhydr) oxides on Cr (VI) reduction by Fe (II)-bearing clay minerals. *Environ. Sci. Technol.* **2019**, 53 (23), 13767–13775.
- (80) Qian, A.; Lu, Y.; Zhang, Y.; Yu, C.; Zhang, P.; Liao, W.; Yao, Y.; Zheng, Y.; Tong, M.; Yuan, S. Mechanistic insight into electron transfer from Fe (II)-bearing clay minerals to Fe (hydr) oxides. *Environ. Sci. Technol.* **2023**, 57 (21), 8015–8025.
- (81) Katz, J. E.; Zhang, X.; Attenkofer, K.; Chapman, K. W.; Frandsen, C.; Zarzycki, P.; Rosso, K. M.; Falcone, R. W.; Waychunas, G. A.; Gilbert, B. Electron small polarons and their mobility in iron (oxyhydr) oxide nanoparticles. *Science* **2012**, 337 (6099), 1200–1203.
- (82) Zhang, Y.; Tong, M.; Lu, Y.; Zhao, F.; Zhang, P.; Wan, Z.; Li, P.; Yuan, S.; Wang, Y.; Kappler, A. Directional long-distance electron transfer from reduced to oxidized zones in the subsurface. *Nat. Commun.* **2024**, 15 (1), 6576.
- (83) Roden, E. E.; Urrutia, M. M. Influence of biogenic Fe (II) on bacterial crystalline Fe (III) oxide reduction. *Geomicrobiol. J.* **2002**, 19 (2), 209–251.
- (84) Liu, C.; Kota, S.; Zachara, J. M.; Fredrickson, J. K.; Brinkman, C. K. Kinetic analysis of the bacterial reduction of goethite. *Environ. Sci. Technol.* **2001**, 35 (12), 2482–2490.

- (85) Kostka, J. E.; Nealson, K. H. Dissolution and reduction of magnetite by bacteria. *Environ. Sci. Technol.* **1995**, 29 (10), 2535–2540.
- (86) Ali, A.; Zafar, H.; Zia, M.; Ul Haq, I.; Phull, A. R.; Ali, J. S.; Hussain, A. Synthesis, characterization, applications, and challenges of iron oxide nanoparticles. *Nanotechnol. Sci. Appl.* **2016**, 9, 49–67.
- (87) Tao, Z.; Zhou, Q.; Zheng, T.; Mo, F.; Ouyang, S. Iron oxide nanoparticles in the soil environment: Adsorption, transformation, and environmental risk. *J. Hazard. Mater.* **2023**, 459, No. 132107.
- (88) Wigginton, N. S.; Haus, K. L.; Hochella, M. F., Jr Aquatic environmental nanoparticles. *J. Environ. Monit.* **2007**, 9 (12), 1306–1316.